

A Systematic Review of Malyugin Rings in Cataract Surgery: Surgical Outcomes, Efficacy and Safety



Salem Abu Al-Burak^{1,#}, Fahad Butt^{1,#}, Jun Bo Wang¹, Emma Anwar², Amit X. Garg^{3,4}, Cindy M. L. Hutnik^{5,6} and Monali S. Malvankar-Mehta^{1,3,5,*}

¹Schulich School of Medicine & Dentistry, Western University, London, ON, Canada

²Faculty of Health Sciences, Western University, London, ON, Canada

³Department of Epidemiology and Biostatistics, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada

⁴Division of Nephrology, Department of Medicine, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada

⁵Department of Ophthalmology, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada

⁶Department of Pathology and Lab Medicine, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada

Abstract:

Introduction: Small pupils complicate cataract surgery and increase intraoperative risk. The Malyugin ring is a widely used mechanical pupil-expansion device lacking a comprehensive evidence synthesis. This review evaluated its efficacy and safety in adults.

Methods: A total of 4 databases (EMBASE, MEDLINE, CINAHL Plus, Web of Science) were searched from inception to October 21, 2024, with an updated search conducted on June 28, 2025. Randomized and observational studies involving adults undergoing cataract surgery with a Malyugin ring were eligible. The outcomes included dilation adequacy, surgical duration, intraoperative complications, endothelial cell loss, and visual acuity. Risk of bias was assessed using RoB 2 and ROBINS-I, and the certainty of evidence was evaluated using GRADE. A narrative synthesis was performed in accordance with PRISMA guidelines (PROSPERO CRD420251069562).

Results: Altogether 10 studies (3,233 eyes) were included. The Malyugin ring provided stable dilation (5.5–7.0 mm) in small pupils and in patients with pseudoexfoliation syndrome. Relative to iris hooks and manual stretching, it was associated with shorter phacoemulsification time (15.2 vs. 19.8 s) and lower endothelial cell loss (9.35% vs. 13.77%). The visual outcomes were favorable, although some studies reported higher rates of iris tears and pupil ovalization compared with those observed with comparator techniques.

Discussion: These findings suggest advantages in surgical efficiency and early visual recovery. However, the evidence comprises 2 randomized and 8 non-randomized studies with risk of bias ranging from low to serious, and no outcomes could be pooled; therefore, comparative conclusions remain tentative.

Conclusion: The Malyugin ring appears effective and generally safe in adult cataract surgery; reported complications were typically not visually significant. Certainty of evidence was low to very low (GRADE), and superiority over alternative devices is not established.

Keywords: Malyugin ring, Pupil expansion device, Small pupil, Cataract surgery, Phacoemulsification, Endothelial cell loss, Systematic review.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Received: May 05, 2026

Revised: July 12, 2026

Accepted: July 15, 2026

Published: August 28, 2026



Send Orders for Reprints to
reprints@benthamscience.net

*Address correspondence to this author at the Schulich School of Medicine & Dentistry, Western University, London, ON, Canada, Department of Epidemiology and Biostatistics, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada and Department of Ophthalmology, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada, Ivey Eye Institute, St. Joseph's Hospital, 268 Grosvenor Street, London, Ontario, N6A 4V2; Tel/Fax: +1 519-685-8500, +1 519-661-3766; ext. 61288; E-mail: Monali.Malvankar@schulich.uwo.ca

*These authors contributed equally to this work.

Cite as: Abu Al-Burak S, Butt F, Wang J, Anwar E, Garg A, Hutnik C, Malvankar-Mehta M. A Systematic Review of Malyugin Rings in Cataract Surgery: Surgical Outcomes, Efficacy and Safety. *Open Ophthalmol J*, 2026; 20: e18743641513562. <http://dx.doi.org/10.2174/0118743641513562260825080417>

1. INTRODUCTION

Cataracts are among the most common causes of remediable blindness, characterized by opacification of the lens. According to World Health Organization data, more than 22 million people in the United States alone had cataracts in 2013 [1, 2]. When visual function is impaired, cataract surgery constitutes one of the most effective treatments available and is associated with improvements in visual acuity, activities of daily living, and reduced mortality [2].

Although cataract surgery is highly effective, it is not without challenges. Specifically, the presence of small pupils (SPs) poses significant difficulty for the procedure. A SP restricts the area within which the surgeon can operate and increases the chances of damaging the pupillary or capsular edge with surgical instruments [3]. Small pupils are also associated with increased intraoperative complications such as posterior capsule rupture (PCR) [4, 5].

There are many methods to achieve adequate pupil dilation, ranging from pharmacologic iris dilation and surgical techniques to mechanical iris expansion. One of the established mechanical dilation devices is the Malyugin ring [6, 7]. It is a single-use, disposable polypropylene ring developed by Dr. Boris Malyugin (Moscow, Russia) and has been reported to offer several advantages. These include reduced trauma to the eye owing to its distributed stretching and gentle holding mechanism, reliable positioning provided by the equidistant loops of the ring, and the absence of any risk of iris aspiration [7]. The Malyugin ring has been reported as an effective tool for managing small pupils in the context of FLACS and intraoperative floppy-iris syndrome [5, 8]. Malyugin rings have also been reported to reduce intraoperative corneal endothelial cell loss, maintain a more stable pupil opening, and offer faster and easier use than iris hooks [9-11]. On the other hand, some studies have reported that Malyugin rings are associated with a higher rate of iris tears than iris hooks and cause significantly greater pupillary distortion in the postoperative period than other mechanical dilation devices, such as the B-HEX ring [3, 12]. Despite the growing body of evidence, there is currently no comprehensive synthesis of data evaluating the outcomes, benefits, and limitations of the Malyugin ring compared with other pupil-dilation methods in cataract surgery.

This systematic review aims to evaluate existing evidence on the use of the Malyugin ring in cataract surgery, providing clinicians and researchers with evidence-based guidance on its optimal use and addressing questions about its efficacy, safety, and comparative performance.

2. METHODS

2.1. Study Design

This systematic review was registered with PROSPERO (registration number: CRD420251069562) and abides by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist [13]. A PICOT framework was developed for the review (**Supplementary Table 1**), with a focus on an adult patient population undergoing cataract surgery with Malyugin rings as a dilating agent.

2.2. Search Strategy

A systematic literature search was conducted in 4 electronic databases: EMBASE (Ovid), MEDLINE (Ovid), CINAHL Plus (EBSCO), and Web of Science, from inception to October 21, 2024. Complete, verbatim search strategies for all 4 databases are provided in **Supplementary Tables 2a-2d**. All records were uploaded to Covidence® for screening. No restrictions on year of publication, study design, or language were applied at the search stage; publication year and language were applied only as eligibility criteria at full-text screening. In addition to the 4 databases, the reference lists of all included studies and relevant reviews were hand-searched for further eligible records (none were identified). The search was updated on June 28, 2025, and no additional eligible studies were identified.

2.3. Study Selection

Studies were eligible if they were written in English, published after 2000, and involved adult patients undergoing cataract surgery in which the Malyugin ring was used as a pupil-expansion device. Eligible designs were randomized controlled trials (RCTs) and observational studies. To be included, studies had to report at least one relevant outcome, such as pupil dilation (measured by pupil diameter), duration of mydriasis, adverse effects (*e.g.*, cardiovascular side effects, increased intraocular

pressure), or intraoperative and postoperative complications.

Studies were excluded if they involved pediatric patients or individuals with contraindications to Malyugin rings or other mydriatic agents. Case reports, commentaries, reviews, and studies without comparison groups were also excluded, as were studies that failed to report relevant outcomes or provided insufficient data for extraction.

Titles and abstracts were screened independently and in duplicate by two reviewers (SA, FB) using Covidence. Full texts were then assessed independently and in duplicate by the same two reviewers, with reasons for exclusion recorded (language, population, study design, intervention, outcome, and date). Disagreements at any stage were resolved by discussion or, where consensus could not be reached, by adjudication from a third reviewer (JW). A PRISMA flow diagram summarizes the study selection process (Fig. 1).




Fig. (1). PRISMA flow diagram demonstrating the identification, screening, and inclusion phases of the systematic review.

2.4. Data Extraction and Synthesis

Data extraction was performed independently by two reviewers (SA, FB). Extracted information included authorship, publication year, country of study, study design, and sample size to provide context for each study. Patient details such as age, gender distribution, and comorbidities were also collected, along with intervention specifics, including dosage, comparator groups, and intervention duration, to facilitate interpretation and applicability of the findings to a broader population.

Data synthesis was conducted narratively, following the Synthesis Without Meta-analysis (SWiM) guideline. Study-level risk-of-bias judgements and the certainty of evidence for each reported outcome, rated using GRADE, are summarized in **Supplementary Tables 3a** and **3b**, respectively. A quantitative meta-analysis was planned a priori but was not performed because no outcome was quantitatively poolable: for each candidate outcome, at most one study reported data in a compatible format (mean and standard deviation, or a 2×2 comparison against a common comparator), while the remaining studies reported medians and interquartile ranges, used non-equivalent comparators, or applied differing devices, sizes, outcome definitions, and follow-up windows (**Supplementary Table 4**). Because synthesis was narrative rather than quantitative, no formal power calculation or minimum information size was applicable. The accumulated sample of 3,233 eyes across 10 studies was nonetheless considered sufficient to support a narrative synthesis, in that it spans the clinically relevant contexts in which the device is used (small pupils, pseudoexfoliation syndrome, intraoperative floppy-iris syndrome, and femtosecond laser-assisted surgery), both commercially available ring sizes (6.25 mm and 7.0 mm), and all principal comparators (iris hooks, B-HEX and OASIS expanders, manual stretching, iris radial cuts, ophthalmic viscosurgical devices, and intracameral mydriatics), with consistent directions of effect within each outcome domain. The accumulated sample, however, was insufficient to permit quantitative pooling or to detect uncommon adverse events, and this is reflected in the GRADE ratings and limitations. All *p*-values reported in this review are those published by the primary studies; no additional statistical testing was undertaken by the review team. Each included study applied the conventional two-sided threshold of $p < 0.05$ to denote statistical significance, and significance is reported here as designated in the source publications. Any discrepancies in data extraction were resolved through discussion and consensus between the reviewers.

2.5. Risk of Bias Assessment

The risk of bias for studies included after full-text screening was assessed independently by two reviewers (SA, EA). For randomized controlled trials (RCTs), the Cochrane Risk of Bias Tool (RoB 2) was used to assess potential biases across selection, performance, detection, attrition, and reporting [14]. For observational studies, the Risk of Bias in Non-Randomized Studies of Interventions

(ROBINS-I) tool was used to assess bias due to confounding, participant selection, intervention classification, deviations from intended interventions, missing data, outcome measurement, and selective reporting [15]. Any disagreements in bias assessments were resolved through discussion and consensus, or by consulting a third reviewer (FB).

3. RESULTS

3.1. Literature Search

A total of 449 publications were identified through comprehensive searches of 4 databases (EMBASE, *n* = 214; MEDLINE, *n* = 87; Web of Science, *n* = 137; CINAHL Plus, *n* = 11). After removing 151 duplicates, 298 records were screened, of which 271 were excluded. Altogether 27 reports were sought and assessed for eligibility, of which 17 were excluded at the full-text stage: 13 evaluated an intervention other than the Malyugin ring or did not provide extractable Malyugin ring-specific outcomes, 3 were not available in English, and 1 involved an ineligible patient population. A total of 10 studies met the inclusion criteria (Fig. 1).

3.2. Study Characteristics

In all, 10 studies (3,233 eyes) evaluated surgical outcomes with the Malyugin ring (Table 1) [3, 5, 8-12, 16-18]. Sample sizes ranged from 21 to 1,426 eyes, with mean ages 57.6–80 years, consistent with typical cataract populations. Most cohorts included both sexes, except Chang (2008), which enrolled only males [8]. Studies were conducted in Europe (*n* = 5), North America (*n* = 3), and Asia (*n* = 2), and were published between 2008 and 2024. Designs comprised 2 randomized trials, 4 prospective non-randomized studies, and 4 retrospective studies, with earlier reports (Chang 2008; Wilczynski *et al.* 2013) providing baseline comparisons. The evidence base thus comprised 2 randomized controlled trials (Gupta *et al.* 2024; Wilczynski *et al.* 2013) and 8 non-randomized studies, 2 of which (Fay *et al.* 2014; Hanna *et al.* 2014) are conference abstracts with limited methodological reporting [8, 9, 12, 17, 18]. This design mix, and the correspondingly mixed risk of bias (low to serious), should be borne in mind when interpreting all comparative findings below.

3.3. Safety and Postoperative Outcomes

Safety outcomes for the Malyugin ring were reported across 6 studies (Table 2). In a retrospective case series from the United Kingdom, Nderitu *et al.* found that the 7.0 mm Malyugin ring was associated with a higher incidence of postoperative inflammation and corneal edema compared to iris hooks [10]. Conversely, Borkenstein *et al.* conducted a prospective study in Austria involving patients with pseudoexfoliation syndrome and found that the 6.25 mm Malyugin ring was well tolerated, with minimal adverse events [16].

In the United States, Fay *et al.* retrospectively assessed changes in endothelial cell density (ECD) and concluded that Malyugin rings caused only minor ECD loss

within safe clinical thresholds [17]. Wilczynski *et al.* similarly found less endothelial cell loss with Malyugin rings than with iris hooks, although more frequent pupil ovalization was observed with the device [9]. In contrast, Balal *et al.* reported that while Malyugin rings improved

postoperative visual acuity, they also demonstrated a higher rate of iris tears relative to iris hooks and intracameral pupil expanders [3]. Hanna *et al.* found that the use of a Malyugin ring was associated with a slightly increased incidence of pseudophakic macular edema (PME) [18].

Table 1. Demographic characteristics of all studies included.

Study	Country	Study Type	Study Group	Sample Size (% females)	Mean Age (SD) Years
Wilczynski <i>et al.</i> (2013) [9]	Poland	Randomized comparative study	Malyugin ring	23 (65.2%)	68 ± 13.8 years
			Manual iris stretching with hooks	17 (70.6%)	73 ± 8 years
Wang <i>et al.</i> (2022) [11]	China	Prospective case-control study	Mechanical manual stretching	16 (62.5%)	66.5 ± 12.1 years
			Iris radial cut	17 (52.9%)	58.2 ± 10.4 years
			Iris-retractor hooks	16 (43.8%)	57.6 ± 11.3 years
			OASIS iris expander	18 (55.6%)	64.9 ± 12 years
			Malyugin ring	16 (56.25%)	66.6 ± 9.0 years
			B-HEX Pupil Expander	16 (56.25%)	59.8 ± 9.0 years
Hanna <i>et al.</i> (2014) [18]	USA	Retrospective review	Malyugin ring	NA	NA
			Iris hooks	NA	NA
Fay <i>et al.</i> (2014) [17]	USA	Retrospective analysis	Malyugin ring	NA	NA
			Cataract Surgery without Malyugin ring	NA	NA
Borkenstein <i>et al.</i> (2018) [16]	Austria	Prospective analysis	Malyugin ring in pseudoexfoliation syndrome (PXF) patients	28 (67.9%)	77.9 ± 7.5 years
Conrad-Hengerer <i>et al.</i> (2013) [5]	Germany	Prospective clinical trial	Epinephrine	850 (50.6%)	73 ± 11 years
			Visco Mydriasis with OVD	850 (50.6%)	73 ± 11 years
			Malyugin ring (7.0 mm)	850 (50.6%)	73 ± 11 years
Chang (2008) [8]	USA	Prospective case series	Malyugin pupil expansion device (5-0 polypropylene)	21 (0%)	76.8 years
Gupta <i>et al.</i> (2024) [12]	India	Randomized control trial	B-HEX pupil expander	64 (46.9%)	70.5 ± 10.12
			Malyugin ring	64 (46.9%)	70.5 ± 10.12
Balal <i>et al.</i> (2021) [3]	United Kingdom	Retrospective case-series	Malyugin ring	1426 (42%)	78.6 years
			Iris hooks	1426 (42%)	78.6 years
			Intracameral phenylephrine	1426 (42%)	78.6 years
Nderitu <i>et al.</i> (2019) [10]	United Kingdom	Retrospective case series	Malyugin ring	135 (43.0%)	80 years
			Iris hooks	45 (47.4%)	80 years

Note: *NA: Not Available.

Table 2. Safety and postoperative outcomes of Malyugin rings as pupil expansion devices in cataract surgery.

Study	Country	Pupil Expansion Device	Device Size	Primary Safety Outcome
Nderitu <i>et al.</i> (2019) [10]	United Kingdom	Malyugin ring/Iris hook	7.0 mm (Malyugin ring)	Higher postoperative inflammation and corneal edema with Malyugin ring
Borkenstein <i>et al.</i> (2018) [16]	Austria	Malyugin ring	6.25 mm	Minimal adverse events and safe for patients with pseudoexfoliation syndrome
Fay <i>et al.</i> (2014) [17]	USA	Malyugin ring	-	Malyugin rings caused minor endothelial cell density (ECD) loss; they remained within safe clinical limits
Wilczynski <i>et al.</i> (2013) [9]	Poland	Malyugin ring/Iris hook	6.0 mm (Malyugin ring)	Improved surgical ease and functional outcomes with less endothelial cell loss than iris hooks; more pupil ovalization
Balal <i>et al.</i> (2021) [3]	United Kingdom	Malyugin ring/Iris hook	6.25/7.0 mm (Malyugin ring)	Malyugin ring was effective in improving VA; showed a higher rate of iris tears
Hanna <i>et al.</i> (2014) [18]	USA	Malyugin ring/Iris hook	-	Malyugin rings achieved faster initial visual recovery; higher incidence of pseudophakic macular edema (PME)

3.4. Efficacy of Pupil Dilation and Maintenance

All 8 studies reporting on the efficacy of pupil dilation supported the Malyugin ring's ability to provide stable, sufficient mydriasis during cataract surgery (Table 3). Nderitu *et al.* found that the 7.0 mm Malyugin ring minimized the number of corneal incisions required during phacoemulsification [10]. Furthermore, in a prospective case series, Chang reported consistent 6.0 mm pupil dilation and reduced intraoperative complications when using a Malyugin pupil expansive device fashioned from 5-0 polypropylene [8].

In a large German clinical trial, Conrad-Hengerer *et al.* demonstrated successful dilation to >5.5 mm in the majority of SP cases using the 7.0 mm Malyugin ring [5]. Fay *et al.* corroborated these findings by noting consistent pupil dilation with only minor ECD loss [17]. Balal *et al.* affirmed the 6.25/7.0 mm Malyugin device's effectiveness in SP cases, demonstrating efficient dilation across a large cohort [3].

Borkenstein *et al.* observed maintained pupil stability in cases of pseudoexfoliation syndrome, with a marked improvement in BCVA postoperatively ($0.48 \log\text{MAR}$ to $-0.05 \pm 0.13 \log\text{MAR}$) [16]. Similarly, Wilczynski *et al.* found that Malyugin ring-treated patients had significantly better postoperative BCVA than those treated with iris hooks (0.75 ± 0.30 vs. 0.56 ± 0.56 ; $p < 0.05$) [9]. Finally, Hanna *et al.* reported faster early visual recovery in the Malyugin group at both postoperative day 1 and week 1 ($\log\text{MAR}$ 0.20 vs. 0.58 at day 1; 0.23 vs. 0.48 at week 1, $p < 0.05$) [18].

3.5. Surgical Performance and Intraoperative Complications

A total of 4 studies assessed the intraoperative

performance of the Malyugin ring (Table 4). In the United Kingdom, Nderitu *et al.* found that the use of the Malyugin ring during phacoemulsification was associated with shorter operative times than with iris hooks, with operative time also varying by surgeon grade (median 20 minutes for consultants vs. 32 minutes for trainees) [10]. Similarly, Wilczynski *et al.* reported a shorter mean phaco time in the Malyugin group compared to iris hooks (15.2 ± 3.9 s vs. 19.8 ± 3.1 s; $p < 0.05$) [9]. In Germany, Conrad-Hengerer *et al.* evaluated the use of the Malyugin ring in femtosecond laser-assisted cataract surgery. They found that the device enabled adequate pupil dilation in 68% of cases with small pupils, without any reported intraoperative complications [5].

Balal *et al.* observed that although intraoperative complication rates were generally comparable across Malyugin rings, iris hooks, and intracameral pupil expanders, the Malyugin group experienced a significantly greater rate of iris tears ($p < 0.05$) [3].

3.6. Comparative Outcomes to Other Pupil Expansion Devices

A total of 7 studies provided comparative analyses of the Malyugin ring and alternative pupil-dilation techniques (Table 5). Nderitu *et al.* reported shorter surgery duration with Malyugin rings; however, iris hooks demonstrated lower postoperative inflammation and edema [10]. Gupta *et al.* found that while both the Malyugin ring and B-HEX pupil expander maintained adequate visualization, the Malyugin ring was associated with significantly more pupillary distortion ($p = 0.029$) [12]. Conrad-Hengerer *et al.* demonstrated superior dilation success with the Malyugin ring relative to viscoadaptive mydriasis using ophthalmic viscosurgical devices (OVDs) or intracameral epinephrine [5].

Table 3. Efficacy of Malyugin rings as a pupil expansion device, investigating the efficacy of pupil dilation and maintenance.

Study	Country	Pupil Expansion Device	Device Size	Primary Efficacy Outcome
Nderitu <i>et al.</i> (2019) [10]	United Kingdom	Malyugin ring/Iris hook	7.0 mm (Malyugin ring)	Malyugin rings reduce corneal incisions required during operation
Chang (2008) [8]	USA	Malyugin pupil expansion device (5-0 polypropylene)	6.0 mm	Maintained consistent 6.0mm pupil dilation and reduced intraoperative complications
Conrad-Hengerer <i>et al.</i> (2013) [5]	Germany	Malyugin ring	7.0 mm	Achieved >5.5 mm pupil dilation; practical control for capsulotomies
Borkenstein <i>et al.</i> (2018) [16]	Austria	Malyugin ring	6.25 mm	Maintain pupil stability in PXF cases; BCVA improved from 0.48 logMAR preop to $-0.05 \pm 0.13 \log\text{MAR}$ post op
Fay <i>et al.</i> (2014) [17]	USA	Malyugin ring	-	Consistent pupil dilation with minor ECD loss
Balal <i>et al.</i> (2021) [3]	United Kingdom	Malyugin ring/Iris hooks	6.25/7.0 mm (Malyugin ring)	Efficient pupil dilation in small pupil cases
Wilczynski <i>et al.</i> (2013) [9]	Poland	Malyugin ring/Iris hook	6.0 mm (Malyugin ring)	Significantly better postoperative BCVA with Malyugin (0.75 ± 0.30 vs. 0.56 ± 0.56 ; $p < 0.05$)
Hanna <i>et al.</i> (2014) [18]	USA	Malyugin ring/Iris hook	-	Faster visual recovery in Malyugin group ($\log\text{MAR}$ 0.20 vs. 0.58 at day 1; 0.23 vs. 0.48 at week 1, $p < 0.05$)

Table 4. Surgical outcomes using Malyugin rings: operating time and complications.


Study	Country	Pupil Expansion Device	Surgical Technique	Primary Surgical Outcome
Nderitu <i>et al.</i> (2019) [10]	United Kingdom	Malyugin ring/Iris hook	Phacoemulsification cataract surgery	Malyugin rings reduced operating times; median operating time: 20 min (consultants) vs. 32 min (trainees)
Balal <i>et al.</i> (2021) [3]	United Kingdom	Malyugin ring/Iris hook/IC PE	Phacoemulsification	Higher rate of iris tears in Malyugin group ($p < 0.05$); overall intraoperative complication rates similar across devices
Wilczynski <i>et al.</i> (2013) [9]	Poland	Malyugin ring/Iris hook	Phacoemulsification	Shorter mean phaco time (15.2 ± 3.9 s vs. 19.8 ± 3.1 s; $p < 0.05$)
Conrad-Hengerer <i>et al.</i> (2013) [5]	Germany	Malyugin ring	Femtosecond laser-assisted cataract surgery with phacoemulsification	Enabled adequate pupil dilation in 68% of small pupil cases; no intraoperative complications occurred

Wang *et al.*, in a six-arm prospective case-control study from China, found that the Malyugin ring achieved good BCVA (median 0.2 logMAR), moderate pupil diameter (median 4 mm), and well-preserved endothelial cell density (median postoperative ECD 1961 cells/mm²), whereas manual iris stretching was associated with the lowest postoperative density (1452 cells/mm²), indicating greater endothelial cell loss; the B-HEX and OASIS devices were less stable or harder to manipulate [11]. Balal *et al.* reported comparable visual acuity improvements across

Malyugin ring, iris hooks, and intracameral phenylephrine, though iris tears were more frequent with the Malyugin ring ($p < 0.05$) [3]. Wilczynski *et al.* found better postoperative BCVA (0.75 vs. 0.56; $p < 0.05$) and lower endothelial cell loss (9.35% vs. 13.77%; $p < 0.05$) with the Malyugin ring than with manual iris stretching [9]. Lastly, Hanna *et al.* reported that the Malyugin group experienced significantly faster visual recovery than the iris hook group during the early postoperative period, although the pseudophakic macular edema rate was slightly higher [18].

Table 5. Comparison of Malyugin ring with alternative pupil dilation techniques in cataract surgery: visual and surgical outcomes.

Study	Country	Comparison Device(s)	Findings of Malyugin Ring	Findings of Comparator
Nderitu <i>et al.</i> (2019) [10]	United Kingdom	Iris hook	Shorter surgery duration	Lower postoperative inflammation and edema
Gupta <i>et al.</i> (2024) [12]	India	B-HEX pupil expander	Maintained sufficient visualization throughout procedure; caused significant pupillary distortion	Less pupillary distortion compared with the Malyugin ring ($p = 0.029$)
Conrad-Hengerer <i>et al.</i> (2013) [5]	Germany	Visco-mydriasis with OVD/epinephrine	Achieved adequate pupil dilation in 68% of small pupil cases; no intraoperative complications	Lower pupil dilation success rates with epinephrine and OVD; some cases required conversion
Wang <i>et al.</i> (2022) [11]	China	B-HEX Pupil Expander/OASIS iris expander/Iris-retractor hooks/Iris radial cut/Mechanical manual stretching	Well-preserved endothelial cell density (median postoperative ECD 1961 cells/mm ²); good BCVA (median 0.2 logMAR); moderate pupil diameter (median 4 mm); simple operation and good stability	Iris hooks similarly preserved endothelial cell density (2114 cells/mm ²); iris radial cut achieved the largest pupil diameter (5.5 mm) but damaged the sphincter; B-HEX less stable; OASIS harder to manipulate; manual stretching had the lowest postoperative density (1452 cells/mm ²), indicating greater endothelial cell loss
Balal <i>et al.</i> (2021) [3]	United Kingdom	Iris hooks/IC PE alone	Similar visual acuity gains across techniques; higher rate of iris tears ($p < 0.05$) in Malyugin group	Iris hooks had higher post-op corneal edema, but not statistically significant; IC PE showed no major complications
Wilczynski <i>et al.</i> (2013) [9]	Poland	Manual iris stretching with hooks	Better postoperative BCVA (0.75 vs. 0.56; $p < 0.05$); lower endothelial cell loss (9.35% vs. 13.77%; $p < 0.05$); easier surgery	Lower postoperative BCVA and greater endothelial cell loss; less stable dilation (4–6 mm)
Hanna <i>et al.</i> (2014) [18]	USA	Iris hook	Faster initial visual recovery (LogMAR 0.20 and 0.23 at day 1 and week 1; $p < 0.05$); PME in 3 cases	Slower initial recovery (LogMAR 0.58 and 0.48 at day 1 and week 1; $p > 0.05$); no PME; similar VA from 1 month on



Study	Risk of bias domains							Overall
	D1	D2	D3	D4	D5	D6	D7	
Nderitu et al. (2018)	-	+	+	+	+	+	+	+
Chang (2008)	X	+	+	+	+	+	-	X
Conrad-Hengerer et al. (2013)	-	+	-	+	+	+	+	-
Borkenstein et al. (2018)	-	+	+	+	+	-	-	-
Fay et al. (2014)	X	+	+	+	+	+	-	X
Wang et al. (2022)	+	+	+	+	-	+	-	+
Balal et al. (2021)	+	+	+	-	+	+	+	+
Hanna et al. (2014)	+	+	-	+	+	+	-	-

Domains:
D1: Bias due to confounding.
D2: Bias due to selection of participants.
D3: Bias in classification of interventions.
D4: Bias due to deviations from intended interventions.
D5: Bias due to missing data.
D6: Bias in measurement of outcomes.
D7: Bias in selection of the reported result.

Judgement
X Serious
- Moderate
+ Low

Fig. (3). Summary of risk of bias assessment using the ROBINS-I Version 2 (2024) tool, evaluating study quality across 7 domains: confounding (D1), classification of interventions (D2), selection of participants into the study (D3), deviations from intended interventions (D4), missing data (D5), measurement of outcomes (D6), and selection of the reported result (D7).

This difference, while statistically significant, does not appear to affect functional outcomes. This aligns with the minor postoperative ovalization observed by Wilczynski *et al.*, which appears to have no lasting clinical impact [9].

The advantages and disadvantages regarding surgical efficiency and pupil shape retention highlight the importance of tailoring device selection to the individual patient's needs and the surgical context. For routine cases or those involving small pupils, the Malyugin ring's efficiency makes it a preferred choice. In contrast, for cases where postoperative pupil aesthetics are a priority, such as in younger patients or those with specific cosmetic concerns, devices like the B-HEX ring may be more suitable.

A plausible device-mechanics explanation for the favorable metrics of the Malyugin ring is its square, four-point engagement of the pupillary margin, which maintains a stable circular aperture under low, evenly distributed tension. Compared with point-loading of individual iris hooks, this distributed hold may reduce localized sphincter microtrauma and iris chafing, which could, in turn, underlie the lower endothelial cell loss and shorter phacoemulsification times reported with the ring. This remains a mechanistic hypothesis rather than a

demonstrated causal pathway, and should be tested directly in adequately powered comparative studies.

Because no outcome was sufficiently comparable to pool quantitatively, subgroup effects are described narratively. By ring size, the 7.0 mm ring was used in larger or more atonic pupils and the 6.25 mm ring in pseudoexfoliation and standard small-pupil cases, with adequate dilation reported across sizes [3, 6, 10, 16]. By surgery type, efficacy was reported in both standard phacoemulsification and femtosecond laser-assisted cataract surgery, where the ring achieved adequate dilation in 68% of small-pupil cases without intraoperative complications [5]. By population, benefit was reported in small pupils, pseudoexfoliation syndrome, and intraoperative floppy-iris syndrome [8]. The protocol pre-specified an adult population and excluded pediatric patients; therefore, evidence-based conclusions regarding pediatric cataract surgery cannot be drawn, and this represents a priority for future research.

Where a higher rate of iris tears with the Malyugin ring was reported, the risk may be mitigated intra-operatively by fully unfolding and centering the ring before manipulation, engaging all 4 scrolls squarely on the pupillary margin under an ophthalmic viscosurgical

device, avoiding over-stretch in fibrotic or pseudo-exfoliative irides, and disengaging and removing the ring before withdrawing the phaco tip [3]. Balanced against this, the direct comparative evidence is mixed: in one study, iris hooks were associated with higher post-operative anterior uveitis and corneal edema than the ring, so device choice should weigh both intraoperative and postoperative complication profiles [10]. Pharmacologic mydriasis remains a complementary first-line strategy: a systematic review of intracameral epinephrine reported effective, well-tolerated dilation during cataract surgery, supporting a stepwise approach in which pharmacologic agents are attempted first, and mechanical expansion is reserved for eyes in which pharmacologic dilation proves inadequate [24].

4.1. Strengths and Limitations

This systematic review provides a structured synthesis of the efficacy and safety of Malyugin rings as a pupil expansion device in cataract surgery. The review followed the PRISMA guidelines and employed a dual-reviewer process for study selection, data extraction, and risk of bias assessment, thereby reducing selection and reporting biases. Risk of bias was rigorously assessed using both the RoB 2 tool for randomized controlled trials and the ROBINS-I tool for observational studies, increasing the credibility and transparency of the findings. Additionally, including studies across a wide range of countries and surgical contexts enhances the external validity of the results. Many of the included studies were prospective in design, allowing for more consistent data collection and minimizing recall bias.

This review has several limitations. Considerable heterogeneity was observed across studies, particularly in terms of surgical technique, device size (6.25 mm vs. 7.0 mm), comparator interventions (*e.g.*, iris hooks, B-HEX, OVDs), and outcome measures such as BCVA, endothelial cell loss, and surgical time. The inclusion of both randomized and non-randomized studies introduced methodological heterogeneity that may affect the strength of our conclusions. Finally, some studies lacked complete reporting of key variables such as gender distribution, follow-up duration, and surgeon experience, which may introduce residual confounding. In addition, 4 additional limitations warrant emphasis. First, no outcome could be quantitatively synthesized: reporting was heterogeneous (means with SD *versus* medians with interquartile ranges), comparators were non-equivalent, and outcome definitions and follow-up windows differed; therefore, only a narrative synthesis was possible, and certainty of evidence was low to very low across outcomes (GRADE). Second, follow-up was short and heterogeneous across studies (predominantly ≤ 1 month, with only a subset reporting to 3–6 months), precluding conclusions about long-term endothelial, refractive, or pupil-shape outcomes. Third, although no language restriction was applied at the search stage, English-language eligibility at the full-text stage resulted in the exclusion of 3 records. As only 3 records were excluded on language grounds, the likely impact on

the conclusions is small; however, language bias cannot be completely excluded. Fourth, 2 included records are conference abstracts with limited methodological detail, further weakening the evidence [17, 18].

CONCLUSION

The Malyugin ring appears to offer advantages in surgical efficiency, safety, and early postoperative outcomes; however, on the predominantly non-randomized evidence available, definitive superiority over alternative expansion devices is not established, and its limitations in certain respects, such as postoperative pupil-shape retention, indicate that device selection should be tailored to the specific needs of each case. Future studies exploring long-term outcomes and broader comparisons with emerging technologies like the B-HEX ring would further refine its role in cataract surgery.

AUTHORS' CONTRIBUTIONS

The authors confirm their contributions to the paper as follows: S.A.A.-B., F.B., C.M.L.H., and M.S.M.-M.: Study conception and design; S.A.A.-B., F.B., J.B.W., and E.A.: Data collection; S.A.A.-B., F.B., A.X.G., C.M.L.H., and M.S.M.-M.: Analysis and interpretation of results; S.A.A.-B. and F.B.: Draft manuscript preparation. All authors reviewed the results, critically revised the manuscript, and approved the final version of the manuscript.

LIST OF ABBREVIATIONS

BCVA	= Best-Corrected Visual Acuity
ECD	= Endothelial Cell Density
ECL	= Endothelial Cell Loss
FLACS	= Femtosecond Laser-Assisted Cataract Surgery
GRADE	= Grading of Recommendations Assessment, Development and Evaluation
IC PE	= Intracameral Phenylephrine
IFIS	= Intraoperative Floppy-Iris Syndrome
IOP	= Intraocular Pressure
logMAR	= Logarithm of the Minimum Angle of Resolution
OVD	= Ophthalmic Viscosurgical Device
PCR	= Posterior Capsule Rupture
PME	= Pseudophakic Macular Edema
PXF	= Pseudoexfoliation Syndrome
RCT	= Randomized Controlled Trial
RoB 2	= Cochrane Risk-of-Bias tool version 2
ROBINS-I	= Risk Of Bias In Non-randomized Studies of Interventions
SP	= Small Pupil
SWiM	= Synthesis Without Meta-analysis

CONSENT FOR PUBLICATION

Not applicable.

STANDARDS OF REPORTING

The study was conducted in accordance with the PRISMA guidelines.

AVAILABILITY OF DATA AND MATERIALS

All data generated or analyzed during this study are included in this published article.

FUNDING

The Pragmatic Trials Training Program, Western University, supported this work.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

SUPPLEMENTARY MATERIAL

The supplementary materials are available on the publisher's website along with the published article.

REFERENCES

- [1] Gollogly HE, Hodge DO, St Sauver JL, Erie JC. Increasing incidence of cataract surgery: Population-based study. *J Cataract Refract Surg* 2013; 39(9): 1383-9. <http://dx.doi.org/10.1016/j.jcrs.2013.03.027> PMID: 23820302
- [2] Moshirfar M, Milner D, Patel BC. *Cataract Surgery*. StatPearls. Treasure Island (FL): StatPearls Publishing 2023.
- [3] Balal S, Jbari AS, Nitiapapand R, *et al.* Management and outcomes of the small pupil in cataract surgery: IRIS hooks, Malyugin ring or phenylephrine? *Eye (Lond)* 2021; 35(10): 2714-8. <http://dx.doi.org/10.1038/s41433-020-01277-0> PMID: 33184489
- [4] Narendran N, Jaycock P, Johnston RL, *et al.* The Cataract National Dataset electronic multicentre audit of 55 567 operations: Risk stratification for posterior capsule rupture and vitreous loss. *Eye (Lond)* 2009; 23(1): 31-7. <http://dx.doi.org/10.1038/sj.eye.6703049> PMID: 18327164
- [5] Conrad-Hengerer I, Hengerer FH, Schultz T, Dick BH. Femtosecond laser-assisted cataract surgery in eyes with a small pupil. *J Cataract Refract Surg* 2013; 39(9): 1314-20. <http://dx.doi.org/10.1016/j.jcrs.2013.05.034> PMID: 23988243
- [6] Sarnicola E, Sarnicola C, Sarnicola V. Dilation devices in cataract surgery. *Curr Opin Ophthalmol* 2023; 34(1): 71-7. <http://dx.doi.org/10.1097/ICU.0000000000000922> PMID: 36484211
- [7] Malyugin B. Small pupil phaco surgery: A new technique. *Ann Ophthalmol (Skokie)* 2007; 39(3): 185-93. <http://dx.doi.org/10.1007/s12009-007-0023-8> PMID: 18025623
- [8] Chang DF. Use of Malyugin pupil expansion device for intraoperative floppy-iris syndrome: Results in 30 consecutive cases. *J Cataract Refract Surg* 2008; 34(5): 835-41. <http://dx.doi.org/10.1016/j.jcrs.2008.01.026> PMID: 18471643
- [9] Wilczynski M, Wierzychowski T, Synder A, Omulecki W. Results of phacoemulsification with Malyugin Ring in comparison with manual iris stretching with hooks in eyes with narrow pupil. *Eur J Ophthalmol* 2013; 23(2): 196-201. <http://dx.doi.org/10.5301/ejo.5000204> PMID: 23112041
- [10] Nderitu P, Ursell P. Iris hooks *versus* a pupil expansion ring: Operating times, complications, and visual acuity outcomes in small pupil cases. *J Cataract Refract Surg* 2019; 45(2): 167-73. <http://dx.doi.org/10.1016/j.jcrs.2018.08.038> PMID: 30527439
- [11] Wang JD, Zhang JS, Li M, Mao YY, Mayinuer Y, Wan XH. Comparison of different pupil dilatation methods for phacoemulsification in eyes with a small pupil. *BMC Ophthalmol* 2022; 22(1): 173. <http://dx.doi.org/10.1186/s12886-022-02402-1> PMID: 35436870
- [12] Gupta S, Shyamsundar K, Pushkar K, *et al.* A randomized control study on post-operative iris distortion following small-pupil cataract surgery using B-HEX pupil expander *versus* Malyugin ring. *Med J Armed Forces India* 2024; 80(5): 560-5. <http://dx.doi.org/10.1016/j.mjafi.2024.05.001> PMID: 39309579
- [13] Page MJ, McKenzie JE, Bossuyt PM, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372(71): n71. <http://dx.doi.org/10.1136/bmj.n71> PMID: 33782057
- [14] Higgins JPT, Altman DG, Gotzsche PC, *et al.* The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; 343: d5928. <http://dx.doi.org/10.1136/bmj.d5928>
- [15] Sterne JAC, Hernán MA, Reeves BC, *et al.* ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016; 355: i4919. <http://dx.doi.org/10.1136/bmj.i4919> PMID: 27733354
- [16] Borkenstein AF, Borkenstein EM. Patient and Surgeon Satisfaction Levels after Using an Acrylic, Hydrophobic, Monofocal IOL and the Malyugin Ring in Pseudoexfoliation Syndrome Patients. *J Ophthalmol* 2018; 2018: 1-5. <http://dx.doi.org/10.1155/2018/3843098> PMID: 29887992
- [17] Fay J, Shrivastava A, Channa P, Madu A. Postoperative course of patients undergoing phacoemulsification cataract extraction with Malyugin Ring™ pupil expansion. *Invest Ophthalmol Vis Sci* 2014; 55(13): 2815-5.
- [18] Hanna J, Kalbag N, Khouri AS. Visual outcomes after uncomplicated complex cataract surgery: Malyugin ring *versus* iris hooks. *Invest Ophthalmol Vis Sci* 2014; 55(13): 2812-2.
- [19] Walkow T, Anders N, Klebe S. Endothelial cell loss after phacoemulsification: Relation to preoperative and intraoperative parameters. *J Cataract Refract Surg* 2000; 26(5): 727-32. [http://dx.doi.org/10.1016/S0886-3350\(99\)00462-9](http://dx.doi.org/10.1016/S0886-3350(99)00462-9) PMID: 10831904
- [20] Lee NSY, Ong K. Risk factors for corneal endothelial cell loss after phacoemulsification. *Taiwan J Ophthalmol* 2024; 14(1): 83-7. <http://dx.doi.org/10.4103/tjo.TJO-D-23-00146> PMID: 38654985
- [21] Almashrafi A, Vanderbloemen L. Quantifying the effect of complications on patient flow, costs and surgical throughputs. *BMC Med Inform Decis Mak* 2016; 16(1): 136. <http://dx.doi.org/10.1186/s12911-016-0372-6> PMID: 27769228
- [22] Cheng H, Clymer JW, Po-Han Chen B, *et al.* Prolonged operative duration is associated with complications: a systematic review and meta-analysis. *J Surg Res* 2018; 229: 134-44. <http://dx.doi.org/10.1016/j.jss.2018.03.022> PMID: 29936980
- [23] Pasquer A, Ducarroz S, Lifante JC, Skinner S, Poncet G, Duclos A. Operating room organization and surgical performance: a systematic review. *Patient Saf Surg* 2024; 18(1): 5. <http://dx.doi.org/10.1186/s13037-023-00388-3> PMID: 38287316
- [24] Butt F, Madani MT, Abu Al-Burak S, *et al.* Efficacy and safety of intracameral epinephrine as mydriatic agent in cataract surgery: A systematic review. *JFO Open Ophthalmology* 2025; 11: 100169. <http://dx.doi.org/10.1016/j.jfop.2025.100169>

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.