



The Open Ophthalmology Journal

Content list available at: <https://openophthalmologyjournal.com>



CASE REPORT

Azzopardi Phenomenon in a Non-neoplastic Phthisical Eye: A Case Report

Wasee Tulvatana^{1,*}, Panitee Luemsamran², Roy Chumdermpadetsuk² and Somboon Keelawat³

¹Department of Ophthalmology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

²Department of Ophthalmology, Rutnin Eye Hospital, Bangkok, Thailand

³Department of Pathology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

Abstract:

Objective:

The Azzopardi phenomenon, known as the deoxyribonucleic acid deposition on various structures due to cellular necrosis, has never been reported in non-neoplastic eyes.

Methods:

We report a case of a 48-year-old man who had congenital nystagmus with poor vision in both eyes, presented with decreased vision and photophobia in his left eye. An exudative retinal detachment was found, which did not respond to systemic steroid treatment. Glaucoma due to occlusio pupillae was later developed. Laser iridotomy and anti-glaucoma medications decreased intraocular pressure to an acceptable level. Vision in the left eye gradually deteriorated during the 10-year clinical course. Evisceration was finally performed due to persistent dull aching ocular pain along with signs of ocular hypotony.

Results:

Histopathological examination showed phthisis bulbi and focal nodular retinal gliosis. The Azzopardi phenomenon was found at the retinal vessel walls, within the retinal layers and along the internal limiting membrane. There was neither evidence of intraocular tumors nor foreign bodies.

Conclusion:

This case demonstrated that the Azzopardi phenomenon could be present in a non-neoplastic eye with a longstanding disease that proceeds to phthisis bulbi.

Keywords: Case report, Azzopardi phenomenon, DNA deposition, Focal nodular retinal gliosis, Phthisis bulbi, Non-neoplastic eye.

Article History

Received: July 28, 2021

Revised: October 14, 2021

Accepted: November 18, 2021

1. INTRODUCTION

The Azzopardi phenomenon, or the Azzopardi effect, was first described by John G. Azzopardi in 1959 [1]. The hematoxyphil (basophilic stained) sheaths around blood vessel walls at the necrotic foci of the oat-cell carcinoma, which was initially believed to be calcification, were first proven by Azzopardi to be the deposition of breakdown products of deoxyribonucleic acid (DNA) liberated from degenerated tumor cells. The material displayed a positive Feulgen reaction and was negative with stains for calcium. He found that almost one-third of oat-cell (small cell) carcinoma specimens had demonstrated this finding.

Most of the reported cases with the Azzopardi phenomenon were from rapidly growing and degenerating tumors, such as pulmonary small cell carcinoma [1, 2], extrapulmonary small cell carcinoma [3 - 5], Burkitt's lymphoma [6], medulloblastoma [7], ependymoblastoma [7], Merkel cell carcinoma [8, 9], primary high-grade neuroendocrine carcinoma of the esophagus [10], squamous cell carcinoma, and polymorphous low grade adenocarcinoma of the oral cavity and the salivary glands [11].

The Azzopardi phenomenon was encountered in a few non-neoplastic cases, *i.e.*, in metal-on-metal arthroplasty cases unrelated to malignancy [12]. This report was later questioned as it was thought to be iron deposition rather than DNA deposition [13]. The phenomenon is also found in functioning parathyroid microcysts, reported in two patients presenting with hyperparathyroidism and hypercalcemia [14]. The

* Address correspondence to this author at the Department of Ophthalmology, Faculty of Medicine, Chulalongkorn University, 1873 Rama 4 Road, Patumwan, Bangkok 10330, Thailand; Tel: +6622564000; E-mail: waseetulvatana@chula.md

proposed mechanisms of these findings were rapid cell cycling or extensive cytoplasmic shrinkage.

Regarding the ophthalmic pathology, Mullaney, in 1968, had originally described this phenomenon in retinoblastoma eyes [15, 16] and later elaborated that the DNA deposition around blood vessels could be found in necrotic areas, calcified foci, cellular areas, and anterior segment of retinoblastoma specimens [17]. Stowe *et al.* in 1979 had studied the incidence of the Azzopardi phenomenon (so-called vascular basophilia) in ocular and orbital tumors. They observed this phenomenon in 6% of retinoblastoma and 1.2% of metastatic carcinoma to the eye and orbit. Oat-cell carcinoma was found to be the major primary cause [18].

As far as we are aware, there have been no previous observations of the Azzopardi phenomenon in a non-neoplastic eye. We herein report this unusual phenomenon in a phthisical eye without evidence of intraocular tumor nor intraocular metal foreign body.

2. CASE PRESENTATION

A 48-year-old male patient, who had a history of congenital nystagmus and poor vision, partially improved with corrective spectacles, was presented to the eye clinic with decreased vision and photophobia in his left eye. He had neither history of ocular trauma nor intraocular surgery. His corrected visual acuity with spectacles (-10.0 D for the right eye and -10.75 D for the left eye) was 20/200 in the right and 5/400 in the left. Intraocular pressure was 21 and 12 mmHg, respectively. Right eye examination appeared to be normal with insignificant retinal pigmentary epithelial changes at the periphery. Left eye examination showed clear cornea, a few cells with flare in the anterior chamber, and exudative retinal detachment. Oral prednisolone 60 mg per day was administered. The retinal detachment did not subside. At a one-month follow up visit, left eye intraocular pressure increased to 38 mmHg from the occlusio pupillae. Iris bombe and high pressure were treated by peripheral laser iridotomy and anti-glaucoma medications. At 6 months, the vision continued to deteriorate and total retinal detachment was observed. At 4 years, the patient had no light perception in the left eye, but the eye was not painful. At 8 years, the intraocular pressure increased again to 35 mmHg. The eye examination showed total occlusion of the pupil, iris neovascularization, localized iris bombe at 2 and 6 o'clock, and the presence of a cyclitic membrane. Cyclophotocoagulation was then planned but was not performed due to well-controlled intraocular pressure after laser iridotomy at the iris bombe areas along with increments of anti-glaucoma medications. The anti-glaucoma eye drops were finally tapered off. At 10 years, the patient reported persistent dull aching pain in the left eye even with the low intraocular pressure (range, 2 to 7 mmHg). The patient agreed to undergo evisceration to relieve the symptoms. Pre-operative physical examination showed no systemic diseases. Complete blood count and chest X-Ray were within normal limits. At a 6-month follow up examination after the operation, the patient was doing well and satisfied with the results. There was no evidence of implant extrusion or other systemic diseases.

3. RESULTS AND DISCUSSION

Histopathological examination of the evisceration specimen showed poorly identifiable structures of the eyeball (Fig. 1A). A neovascular membrane at the anterior surface of the iris, partially calcified lens, and a few small lymphoid aggregates in the choroid were observed. The retina was disorganized, and it showed a localized glial mass representing fibrous metaplasia, occupying approximately one-fifth of the vitreous cavity (Fig. 1B) along with small foci of osseous metaplasia (Fig. 1C) and multinucleated giant cells in the vicinity of numerous cholesterol clefts (Fig. 1D). Deep basophilic staining encircling the retinal vascular channel walls as well as scattered small basophilic clumps of varying sizes were widespread among the poorly identifiable retinal layers (Fig. 1E). Linear deep basophilic staining of the internal limiting membrane was also found (Fig. 1F). The specimen did not show any viable or necrotic tumor cells. There was no evidence of intraocular foreign bodies. The intraretinal basophilic clumps (Fig. 2A) were negative by von Kossa stain for calcium (Fig. 2B). Perl's Prussian Blue stain for iron was negative. The final pathological diagnosis was phthisis bulbi with focal nodular retinal gliosis [19], and the Azzopardi phenomenon at the posterior part of the eyeball, *i.e.*, around the retinal vessels, within the retinal layers, and along the internal limiting membrane.




Fig. (1). Histopathology of the lesion. Photomicrographs of hematoxylin and eosin stains demonstrating (A) the lens with surrounding cyclitic membrane and poorly identifiable other intraocular structures ($\times 4$ power); (B) a localized glial mass occupying approximately one-fifth of the vitreous volume ($\times 10$ power); (C) osseous metaplasia ($\times 40$ power); (D) numerous cholesterol clefts admixed with multi-nucleated giant cells ($\times 40$ power); (E) the Azzopardi phenomenon, as deep basophilic staining around the retinal vessel walls, intra-retinal clumps ($\times 40$ power); and (F) along the internal limiting membrane ($\times 40$ power).




Fig. (2). Histopathology of the retina. Photomicrograph of hematoxylin and eosin stains (A) demonstrating intraretinal basophilic clumps ($\times 40$ power); (B) and the identical area which is negative by von Kossa stain for calcium ($\times 40$ power).

Although the Azzopardi phenomenon is not unexpected in rapidly growing tumor, this phenomenon is extremely rare in non-neoplastic cases. Reports in non-neoplastic lesions showed this effect in only a few cases with metal-to-metal interaction and rapidly growing cells in functioning parathyroid microcysts [12 - 14].

We searched for the literature using the keywords “Azzopardi phenomenon” OR “Azzopardi effect” in PubMed, Scopus, and Google Scholar databases; there has been no previous report of this phenomenon in non-neoplastic eyes.

The presence of cyclitic membrane, lens calcification, focal nodular retinal gliosis, and cholesterol clefts represented a prolonged course of the disease. We did not find any evidence of tumor cells or intraocular foreign bodies in this eye.

CONCLUSION

This case report is an example of the Azzopardi phenomenon in longstanding ophthalmic pathology with evidence of inflammation and possibly from extensive cellular necrosis during the prolonged clinical course of the disease. We, therefore, conclude that the Azzopardi phenomenon can be present in non-neoplastic eye that proceeds to phthisis bulbi.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The research ethics review was exempted by the Institutional Review Board of the Faculty of Medicine, Chulalongkorn University (Certificate of exemption number 001/2021).

HUMAN AND ANIMAL RIGHTS

The report adhered to the tenets of the Declaration of Helsinki as amended in 2008.

CONSENT FOR PUBLICATION

Written informed consent was obtained from the patient.

AVAILABILITY OF DATA AND MATERIALS

The authors confirm that the data supporting the findings of this study are available within the article, and further information is available from the corresponding author, [W.T.], on special request.

STANDARDS OF REPORTING

CARE guidelines have been followed in this study.

FUNDING

None.

CONFLICTS OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

We thank Ms. Sranya Phaisawang and the English editing service, Research Affairs, Faculty of Medicine, Chulalongkorn University, for assistance in the manuscript.

REFERENCES

- [1] Azzopardi JG. Oat-cell carcinoma of the bronchus. *J Pathol Bacteriol* 1959; 78: 513-9. [http://dx.doi.org/10.1002/path.1700780218] [PMID: 13795444]
- [2] Farchoukh L, Pantanowitz L, Xing J, Monaco SE. Azzopardi phenomenon associated with small cell carcinoma. *Diagn Cytopathol* 2019; 47(8): 837-8. [http://dx.doi.org/10.1002/dc.24188] [PMID: 30963714]
- [3] Pritt BS, Cooper K. The azzopardi phenomenon. *Arch Pathol Lab Med* 2003; 127(9): 1231. [http://dx.doi.org/10.5858/2003-127-1231-TAP] [PMID: 12951999]
- [4] Dolzhansky OV, Kaabak MM, Solovyeva SE, Fedorov DN, Pigolkin YI. Disseminated combined small cell carcinoma with graft involvement after kidney transplantation from a deceased donor to two patients. *Arkh Patol* 2017; 79(6): 53-9. [http://dx.doi.org/10.17116/ptol201779653-59] [PMID: 29265078]
- [5] Miranda-Aquino T, Pérez-Topete SE, Hernández-Portales JA, Casillas-Romo A, López-Chuken Y. Gastroesophageal junction small cell carcinoma. *Med Interna Mex* 2017; 33(3): 403-6.
- [6] Peng SL, Cheng CN, Chang KC. Burkitt lymphoma with azzopardi phenomenon. *Arch Pathol Lab Med* 2007; 131(5): 682-3. [http://dx.doi.org/10.5858/2007-131-682-BLWAP] [PMID: 17488151]
- [7] Takei H, Adesina AM, Bhattacharjee MB. The azzopardi phenomenon. *Arch Pathol Lab Med* 2007; 131(4): 518. [http://dx.doi.org/10.5858/2007-131-518-TAP] [PMID: 17425376]
- [8] Vazmitel M, Michal M, Kazakov DV. Merkel cell carcinoma and azzopardi phenomenon. *Am J Dermatopathol* 2007; 29(3): 314-5. [http://dx.doi.org/10.1097/DAD.0b013e318033901b] [PMID: 17519638]
- [9] Vazmitel M, Michal M, Shelekhova KV, Sima R, Mukensnabl P, Kazakov DV. Vascular changes in merkel cell carcinoma based on a histopathological study of 92 cases. *Am J Dermatopathol* 2008; 30(2): 106-11. [http://dx.doi.org/10.1097/DAD.0b013e318162eaa2] [PMID: 18360111]
- [10] Huang Q, Wu H, Nie L, et al. Primary high-grade neuroendocrine carcinoma of the esophagus: A clinicopathologic and immunohistochemical study of 42 resection cases. *Am J Surg Pathol* 2013; 37(4): 467-83. [http://dx.doi.org/10.1097/PAS.0b013e31826d2639] [PMID: 23426118]
- [11] Chatterjee K. The azzopardi phenomenon. *J Oral Maxillofac Pathol* 2015; 19(3): 280-1. [http://dx.doi.org/10.4103/0973-029X.174619] [PMID: 26980952]
- [12] Zustin J, Skinner JA, Hart AJ. Azzopardi phenomenon in cystic pseudotumours associated with retrieved metal-on-metal arthroplasty. *Hum Pathol* 2016; 51: 134-7. [http://dx.doi.org/10.1016/j.humpath.2016.01.009] [PMID: 27067791]
- [13] Natsu S, Langton DJ, Sidaginamale RP. Azzopardi phenomenon reported in metal-on-metal arthroplasties is in fact iron encrustation of blood vessels. *Hum Pathol* 2017; 62: 245-6. [http://dx.doi.org/10.1016/j.humpath.2016.09.033] [PMID: 27864118]
- [14] Handra-Luca A, Tissier F. Infracentimetric parathyroid cysts in hyperparathyroidism. *Pathol Res Pract* 2018; 214(3): 455-8. [http://dx.doi.org/10.1016/j.prp.2017.09.018] [PMID: 29496307]

- [15] Mullaney J. D.N.A. in retinoblastoma. *Lancet* 1968; 2(7574): 918. [http://dx.doi.org/10.1016/S0140-6736(68)91091-X] [PMID: 4176171]
- [16] Mullaney J. Retinoblastomas with DNA precipitation. *Arch Ophthalmol* 1969; 82(4): 454-6. [http://dx.doi.org/10.1001/archophth.1969.00990020456005] [PMID: 4186574]
- [17] Mullaney J. Retinoblastoma, a review and some new aspects. *Ir J Med Sci* 1969; 8(2): 57-64. [http://dx.doi.org/10.1007/BF02958908] [PMID: 4892363]
- [18] Stowe GC III, Zakov ZN, Albert DM, Smith TR, Sang DN, Craft JL. Vascular basophilia in ocular and orbital tumors. *Invest Ophthalmol Vis Sci* 1979; 18(10): 1068-75. [PMID: 225286]
- [19] Jakobiec FA, Thanos A, Stagner AM, Grossniklaus HE, Proia AD. So-called massive retinal gliosis: A critical review and reappraisal. *Surv Ophthalmol* 2016; 61(3): 339-56. [http://dx.doi.org/10.1016/j.survophthal.2015.12.002] [PMID: 26703887]

© 2021 Tulvatana *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.