

# The Prevalence of Contact Lens Wear and Improper Use Among Students of Al-Baha University, Saudi Arabia



Mahadi Bashir<sup>1,2,\*</sup>, Ali Hendi Alghamdi<sup>1</sup>, Aimun A. E. Ahmed<sup>3,4</sup>, Sarah S. Taishan<sup>5</sup>, Wafaa S. Taishan<sup>5</sup>, Mohammed G. Alghamdi<sup>5</sup>, Hind A. Hendi Alghamdi<sup>5</sup>, Lara M. Alsakka<sup>5</sup>, Njood A. Alghamdi<sup>5</sup>, Hadeel M. Alzahrani<sup>5</sup> and Asail S. Alghamdi<sup>5</sup>

<sup>1</sup>Unit of Ophthalmology, Department of Surgery, Faculty of Medicine, Al-Baha University, Al-Baha 23334, KSA

<sup>2</sup>Department of Surgery, Faculty of Medicine, Bahry University, Alkadroo, Sudan

<sup>3</sup>Department of Pharmacology, Faculty of Medicine, Al-Baha University, Al-Baha 23334, KSA

<sup>4</sup>Department of Pharmacology, Faculty of Pharmacy, Omdurman Islamic University, Omdurman, Sudan

<sup>5</sup>Faculty of Medicine, Al-Baha University, Al-Baha 23334, Saudi Arabia

## Abstract:

**Background:** This study aims to evaluate the knowledge and practice of contact lens wear (CLW) among Al-Baha University medical students.

**Methods:** A cross-sectional study was carried out using an online self-administered questionnaire via Google Forms. The questionnaire had three domains: sociodemographic characteristics of the participants, CLW hygiene practices, and CLW-related complications.

**Results:** A total of 432 students were enrolled. Almost half of them (203,47%) were CLs wearers. Almost half of the participants, 95 (46.8%), used CL for therapeutic and 108 (53.2%) cosmetic purposes. 81 (40%) wore their CLs for a duration of (1-10 hours/day), 149 (73.4%) did not sleep while wearing their CLs, 61 (30%) shared theirs with others, 110 (54.2%) washed their hands before putting on their CLs and 90 (44.2%) replaced their CL cleaning solution regularly. Only 10 (5%) of the participants did not experience any complications related to CLW, while 108(52.6%), 102 (50%), and 97 (47.8%) experienced dry eye, tears, and foreign body sensations, respectively. Moderate satisfaction of the users was reported.

**Conclusion:** The medical students at Al-Baha University showed a high CLW ratio and used CL predominantly for cosmetic reasons; both male and female students were very aware of hygienic practices, and moderate satisfaction, with moderate complications, was reported.

**Keywords:** Contact lens, Prevalence, Improper use complications, Medical students, Al-Baha University, Eye glasses.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

\*Address correspondence to this author at the Unit of Ophthalmology, Department of Surgery, Faculty of Medicine, Al-Baha University, Al-Baha 23334, KSA; Tel/Fax: +9667 7274111, +966 7 7247272; E-mail: Kocap72@gmail.com

Cite as: Bashir M, Alghamdi A, Ahmed A, Taishan S, Taishan W, Alghamdi M, Alghamdi H, Alsakka L, Alghamdi N, Alzahrani H, Alghamdi A. The Prevalence of Contact Lens Wear and Improper Use Among Students of Al-Baha University, Saudi Arabia. *Open Ophthalmol J*, 2024; 18: e18743641306553. <http://dx.doi.org/10.2174/0118743641306553240313061238>



Received: February 03, 2024

Revised: February 29, 2024

Accepted: March 06, 2024

Published: ?? ??, 2024



Send Orders for Reprints to [reprints@benthamscience.net](mailto:reprints@benthamscience.net)

## 1. INTRODUCTION

Recently, there has been a progressive expansion in the wear and demand of contact lenses (CLs) due to

significant developments in the field of contactology [1] [2]. CLs are more effective than eyeglasses in treating vision complaints. In addition, they do not change the

image size on the retina, so anisometropia correction is unaffected [3, 4]. CL use leads to greater satisfaction with refractive error correction [5]. CLs are classified as hard, rigid gas-permeable, or soft lenses. Cosmetic CLs are used for non-medical problems [6], and CLs can be classified as therapeutic, cosmetic, optical diagnostic, operative, preventive, and occupational [7, 8] also based on manufacturing material, CLs are classified into Filcons and Facons [7]. The efficacy of CLs can be affected by refractive index, oxygen permeability, wettability, and water content, besides Light transmission, heat resistance, dimensions, and flexural stability [9-11]. CLW presents various potential threats and can cause complications [12, 13], adverse effects of CLW mostly affect conjunctiva and cornea [8, 14]. In addition to the potential for discomfort, wearing CL may carry the risk of infections [6, 14].

Corneal infection or microbial keratitis is the most serious complication of CLW and can lead to blindness [14, 15]. Recent studies confirmed that pathogenic microorganisms adhere to CLs and survive in lens solutions [16]. Fortunately, recent advances in CL material and shape design, along with strict adherence to regular replacement and hygienic care protocols, have significantly reduced discomfort complaints and improved compliance with CLW [17]. However, CL manufacturing still faces many difficulties because of CLW problems [18].

Numerous potentially pathogenic and nonpathogenic organisms make up the microbiological lens pollutants [16]. According to research done in the Kingdom of Saudi Arabia (KSA) a high percentage of CLs were sold in regular stores without prescriptions [19]. A United States study also found that CL wearers are more likely to experience symptoms such as burning, itching, or tearing of the eyes [5]. Furthermore, the lack of growth in CL use in Europe has been attributed in part to the discontinuation of CL wear supply. Additionally, studies have been done on wearers' satisfaction and retention [17, 20-22]. The literature also discusses how various environmental conditions, including temperature and humidity, affect CL wearers' tear films [23, 24]; for this reason, CLs are now being used in ocular surface disease, leading to surface hydration, protection from environmental effects and injury from abnormal lids as well as drug provision to the ocular surface [25].

Studies have found that the prevalence of CLW complaints was higher in soft CLW [26], most of whom have ocular signs and symptoms. Nonetheless, the number of complaints lowered when soft CL wearers are refitted with new-generation silicone hydrogel CLs [26].

Many studies focus on and examine adult CL wearers' eye complaints, such as dryness [12, 14]. The literature also covered the short-term physiological effects of refitting [27, 28], as well as the symptoms, signs, and duration of discomfort experienced by CLs wearers [3, 29]. As shown in a review, soft CLW has also improved [2, 29] [30]. Despite an increase in CLW among medical students

at Al-Baha University, lack of knowledge and improper use were seen; therefore, it is crucial to assess the status of CLW. This study intends to evaluate CLW-related knowledge, current use status, pattern, and improper usage among medical students at Al-Baha University and to report any complications.

## 2. MATERIALS AND METHODS

A cross-sectional study was carried out in Saudi Arabia's Al-Baha area from January 2022 to November 2023. Study participants were recruited from Al-Baha University; they agreed to take part, were of both sexes, could read, and had a social media account. The sample size was calculated using the Epi Info™ (A public domain suite of interoperable software tools) based on a 95% confidence interval, a 5% margin of error, and the total number of Al-Baha University students. The estimated sample size was 385, but it was adjusted to 432 to compensate for a 10% non-response. This study employed an online self-administered questionnaire via Google Forms. The generated link was shared through social networks (*i.e.*, WhatsApp and Telegram). The goal of this research was explained in the interface. A validated questionnaire based on earlier studies was used. The questionnaire had three domains. The first included the participants' sociodemographic characteristics, such as age group, gender, nationality, and residence. The second consisted of questions about CL hygiene practices, while the third centered on CLW-related complications.

### 2.1. Statistical Analysis


The collected data were analyzed using SPSS Version 25, and the variables were then reviewed and presented using descriptive statistics. The categorical variables such as gender, age group, and nationality were summarised and reported in terms of frequency distribution. The reliability test was evaluated using the Cronbach alpha value, which ranged from 0 to 1, and the estimated value was 0.6 - 0.7. The chi-square test was used to evaluate and quantify the associations between sociodemographic factors and CLW. Statistical significance was set at  $p \leq 0.05$ .

### 2.2. Ethical Considerations

The Al-Baha University, Faculty of Medicine Research & Ethics Committee approved this study, which was given the approval number: REC/SUR/BU-FM/2022/43. Our institution committee checks the fulfillment rules of the Helsinki Declaration for involving human subjects in the research. Before enrolling in the study, informed consent was obtained from each participant.

## 3. RESULTS

A total of 432 students were enrolled in the study, 68.6% ( $n=210$ ) of the students' age ranged from 20-22 years. 68.5% ( $n=296$ ) of the population were women. Of the 432, 203 (47%) were contact lens users (Fig. 1) other sociodemographic characteristics are shown in Table 1.



**Fig. (1).** Percentage of CL use.

**Table 1. Sociodemographic characteristics among Al-Baha University students, N=432.**

Variable		Number	(%)
Gender	Male	136	31.5
	Female	296	68.5
Age group (Years)	18-20	117	27.1
	20-22	210	48.6
	23-25	97	22.5
	26-30	2	0.5
	More than 30	6	1.4
Nationality	Saudi Arabian	427	98.8
	Not Saudi Arabian	5	1.2
Marital status	Single	401	92.8
	Married	29	6.7
	Divorced	2	0.5
City	Al-Baha	266	61.6
	Al Aqiq	74	17.1
	Tehama	53	12.3
	Others	39	9

176 of 203 (86.7%) are women. It was found that 46.8% (n=95) and 53.2% (n=108) of the participants used CLs for therapeutic and cosmetics, respectively. The educational levels are shown in Fig. (2). Regarding hygiene practice, we found that 51.6% (n=105) of students use their contact lens for more than 6 hours, 73.4% (n=149) never sleep with the lens, and the results revealed that 93.8% (n=190) of them wash their hands before putting on the lens, and 92% (n=187) replace the cleaning solution. About 30% (n=61) of the participants share the lens with others as shown in Table 2. Regarding the complication related to the use of contact lenses, the results showed that only 5% (n=10) of the participants

experienced any complications, whereas 52.6% (n=108), 50% (n=102), 47.8% (n=97) reported dry eye, tears, and foreign body sensation, respectively (Fig. 3).

Table 3 shows the feedback experience of the users; 58.6% (n=119) found it useful and 53.7% (n=109) changed their life. The Chi-square test was conducted to estimate the association between sociodemographics and contact lenses; the results reported that females use frequently than males. There was a statistically significant association,  $p = 0.0001$ . A higher percentage of women 86.7% (n=176) use CL compared to men 13.3% (n=27). And when applied to the age group, it was also statistically significant with  $p$  value =0.02 (Table 4).



(Table 2) contd....

Hygienic Practices		Number	(%)
Sharing lenses with friends	Never	142	70
	Rare	19	9.4
	Sometimes	27	13.3
	Much	9	4.3
	Always	6	3
Do you change the lens if you notice changes in it?	Never	12	6
	Rare	8	4
	Sometimes	44	21.7
	Much	30	14.7
	Always	109	53.7




Fig. (3). Reported CL-related complications.

Table 3. Feedback on CL experience among Al-Baha University students, N (%).

Experience Feedback		Number	(%)
Is it useful in your life?	Useful	119	58.6
	Maybe	76	37.4
	It is not useful	8	4
Is it expensive?	Expensive	109	53.7
	Neutral	84	41.3
	Cheap	10	5
Is contact lenses a groundbreaking innovation in ophthalmology?	Yes	138	68
	No	5	2.5
	Maybe	60	29.6

**Table 4. Relationship between CL use and population variables.**

Pattern		Users (n=203)	No-users(n=229)	X <sup>2</sup> (p-value)
Gender	Male	27(13.3)	109(47.6)	58.7(0.001) ***
	Female	176(86.7)	120(52.4)	
Age group(Y)	18-20	68(23.6)	49(21.4)	11.6 (0.02) **
	21-22	86(42.4)	124(54.1)	
	23-25	45(22.2)	52(22.7)	
	26-30	0(0.0)	2(0.9)	
	More than 30	4(1.9)	2(0.9)	

Note: X2 chi-square test., \*\*\*, p≤0.001; \*\*, p≤0.01; ns, not statistically significant.

**Table 5. Relationship between gender and the purpose of CL use.**

		Use		Chi-square (p-value)
		Therapeutic	Cosmetic, N (%)	
Gender	Male, N (%)	20(21)	7(6.4)	63.7 (0.0001)
	Female, N (%)	75(79)	101(93.6)	
Total		95	108	-

**Table 6. Reliability test result.**

Scale	Items	Cronbach's Alpha	ICC (95%CI)	Value P value
Hygiene	5	0.613	0.613(0.522-0.691)	0.000<

The Chi-square test was conducted to assess the relationship between gender and use, which showed a statistically significant value that 93.6% (n=101) women use contact lenses for cosmetic use more than males (p value= 0.0001) (Table 5). The questionnaire's reliability was measured by Cronbach's acceptable alpha coefficient. And the consistency between classes is 0.613(0.522-0.691) with a p-value <0.001 (Table 6).

#### 4. DISCUSSION

In this study, all levels of medical students took part with various percentages; this result is worthy evidence for sample coverage [31]. Moreover, an acceptable alpha Cronbach's value > 0.70 indicates satisfactory internal consistency for a scale [32]. On the other hand, the prevalence of CL wear in this study was 53%, this rate is higher than the rate of medical students of King Faisal University (KFU), Al Ahsa, KSA, 2021, which was 47.1% [30], and students of Umm Al-Qura University (UQU) Makkah, KSA, 2016, (50.1%) [29], and study among Saudi female students (47%), 2020 [33], and significantly higher than medical students of King Abdul-Aziz University (KAU), Jeddah, KSA, 2018, which reported prevalence of CL wear to be 40% [2].

Strangely, in a study conducted in Chengdu, China, the rate of CL wear was only 20%, which is much lower than this study [34]. Reversely, a study in Riyadh, KSA, 2014, assessed CL wearing among university female students and reported a prevalence of 70.2% [19], as far as we know, this is the highest rate of CL wear in KSA. The reason for the wide difference in results between Saudi

universities may be due to differences in times of studies or populations studied [2], but worldwide, the rate of CL wear among medical students is widely different [15, 35].

From another view, the majority of CL wearers in this study were females (86.7%) this is consistent with the study conducted among medical students of (KAU), Jeddah, and KSA, where females reported a prevalence of 95% [2], and similarly in a study among medical students of (KFU), Al Ahsa, KSA, reported 63.3% [30], and same like a study in Jordan, 2020, conducted among the university population found that 89% of the CL wearers were female [36], moreover in China, a study in eight different universities reported that (82.15%) of CL wearer were female [35], and among medical students of University of Malaya, 87.6% of CL wearer were female [37].

The study revealed that Cosmetic use was the main purpose of CLW among females 93.6% compared to males (6.4%). The Chi-square test revealed a statistically significant value concerning cosmetic CL wear among females more than males (p value= 0.0001); this result is supported by other similar studies conducted inside KSA, which reported high rates of CL wear for cosmetic purposes among the university female students [2, 19, 29], and also studies conducted outside KSA [37-39].

Concerning CL hygienic practices, almost half of the participants responded to the questions, and the majority of them were female; this is logical because females represent 68.5% of the studied population and 86.7% of CL wearers, and this is also in favor of other published

studies [2, 30, 34, 36, 40], and happily the results were highly satisfactory regarding the time of CL wearing per day (74%), sleeping with CL (73.4%), never sharing CL with others (70%), washing hands before putting on CL (75%), and replacing the cleaning solution (66.4%). These results reflect an elevated level of knowledge, practice, and compliance of Al-Baha medical students concerning hygienic guidelines for CL wear, in comparison, worldwide and local studies have revealed different results of complications and compliance of CLW [30, 40-42] [42].

Results of this study showed that; dryness, tearing, foreign body sensation, and sensitivity to light, are the main complaints reported among CL wearers of Al Baha medical students, while infections are the least one, and although keratitis in this study is the least complication, it is the worst and can lead to severe vision morbidity [6, 11, 42, 43], possibly the low prevalence of infections, including keratitis, among CL wearers of Al Baha medical students may be due to their close adherence to the hygienic protocol, because a lot of studies discovered strong relation between hygiene behaviors of CL wearers and eye complications [16, 42], this is including types of CLs and period of use [44], this is also supported by study conducted in the United States which found that microbial keratitis and dryness were frequently associated with extended CL wearer leading to frequent visiting eye emergency units [43], moreover it led to change in, ocular morphology, corneal sensitivity, and tear film [12]. In the present study, conjunctivitis represented 24% of complaints, but they need more care when they use CLs [13]. Concerning the satisfaction of students, about 60% of the contributors responded positively regarding the benefit of CLs wearing, and 68% looked at CL as an innovative tool in ophthalmology while more than half thought that CLs are expensive. In contrast to these results, a study assessed the satisfaction of CLs wear revealed that significant part of CL users was not satisfied and liable to discontinuation [22, 23], similarly, another study reported that 23% of the wearers had permanently stopped CL use mainly due to dryness and discomfort [21], and another study concluded that a large number of lapsed CL users can be effectively refitted with CLs [17]. A study found that the dropout of new lens wearers was 74%, with many laps in the first two months, and, in addition to lens power, material, and type, the frequent purchase are the reasons [17, 19].

#### 4.1. Limitations of Study

The limitations of this study are that it is an online survey with a limited time.

#### CONCLUSION

Female medical students from Al-Baha University showed a high CL wear ratio, and both male and female students were highly aware of hygienic practices. Females used CLs predominantly for cosmetic reasons, unlike male students, who use them mainly for therapeutic purposes. On the other hand, a moderate satisfaction level with moderate complications, such as dryness, tearing, and foreign body sensation, was observed among study

participants. Although a low ratio of Keratitis was reported, it is considered as a serious complication. Still small ratio of students had malpractice in CL wearing and needed education.

#### LIST OF ABBREVIATIONS

CLW	=	Contact Lens Wear
CL	=	Contact Lens
KSA	=	Kingdom of Saudi Arabia
KAU	=	King Abdul-Aziz University
KFU	=	King Faisal University
UQU	=	Umm Al-Qura University

#### ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The Al-Baha University, Faculty of Medicine Research & Ethics Committee approved this study, which was given the approval number: REC/SUR/BU-FM/2022/43.

#### HUMAN AND ANIMAL RIGHTS

No animal were used that are the basis of this study. Our institution committee checked the fulfilment rules of the Helsinki Declaration for involving human subjects in the research.

#### CONSENT FOR PUBLICATION

Before enrolling in the study, informed consent was obtained from each participant.

#### STANDARDS OF REPORTING

STROBE guidelines were followed.

#### AVAILABILITY OF DATA AND MATERIALS

The data and supportive information are available within the article.

#### FUNDING

None.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

#### ACKNOWLEDGEMENTS

Declared none.

#### REFERENCES

- [1] Ferreiro SAV, Bellido ML. Historical evolution of contact lenses. Arch Soc Esp Oftalmol 2012; 87(8): 265-6. <http://dx.doi.org/10.1016/j.oftal.2012.04.009> PMID: 22794177
- [2] Ibrahim NKR, Seraj H, Khan R, Baabdullah M, Reda L. Prevalence, habits and outcomes of using contact lenses among medical students. Pak J Med Sci 2018; 34(6): 1429-34. <http://dx.doi.org/10.12669/pjms.346.16260> PMID: 30559798
- [3] Pillay R, Hansraj R, Rampersad N. Historical development, applications and advances in materials used in spectacle lenses and contact lenses [Response to Letter]. Clin Optom 2020; 12: 201-2. <http://dx.doi.org/10.2147/OPTO.S289792> PMID: 33204202

- [4] Mandell RB, Harris M, Polse KA. History of contact lenses at the University of California, Berkeley, school of Optometry. *J Am Optom Assoc* 1998; 69(6): 376-86. PMID: 9646584
- [5] Walline JJ, Gaume A, Jones LA, et al. Benefits of contact lens wear for children and teens. *Eye Contact Lens* 2007; 33(6): 317-21. <http://dx.doi.org/10.1097/ICL.0b013e31804f80fb> PMID: 17993828
- [6] Ahuja M. Contact lens wear and microbial keratitis. *J Indian Med Assoc* 2002; 100(11): 664-6. PMID: 12797639
- [7] Yuan W, Zhao F, Liu X, Xu J. Development of corneal contact lens materials and current clinical application of contact lenses: A review. *Biointerphases* 2023; 18(5): 050801. <http://dx.doi.org/10.1116/6.0002618> PMID: 37756594
- [8] Bhanot K, Jefferys S, Clipstone K, Guest S, Blanch RJ. Contact lens-related complications in austere conditions among military personnel: A systematic review. *BMJ Mil Health* 2023; 12: e002476. <http://dx.doi.org/10.1136/military-2023-002476> PMID: 37699733
- [9] Costa D, De Matteis V, Treso F, et al. Impact of the physical properties of contact lens materials on the discomfort: Role of the coefficient of friction. *Colloids Surf B Biointerfaces* 2024; 233: 113630. <http://dx.doi.org/10.1016/j.colsurfb.2023.113630> PMID: 37956592
- [10] Contact Lenses. 2023. Available from: <https://clspectrum.com/issues/2024/januaryfebruary/contact-lenses-2023/>
- [11] Lee SE, Kim SR, Park M. Oxygen permeability of soft contact lenses in different pH, osmolality and buffering solution. *Int J Ophthalmol* 2015; 8(5): 1037-42. <http://dx.doi.org/10.3980/j.issn.2222-3959.2015.05.33> PMID: 26558223
- [12] Tena SMÁ, Perez MC, Collar VC, Peregrina AC. Long-term effect of contact lens wear: A citation network study. *Cont Lens Anterior Eye* 2022; 45(1): 101527. <http://dx.doi.org/10.1016/j.clae.2021.101527> PMID: 34732300
- [13] Urgacz A, Mrukwa E, Gawlik R. Adverse events in allergy sufferers wearing contact lenses. *Postepy Dermatol Alergol* 2015; 3(3): 204-9. <http://dx.doi.org/10.5114/pdia.2015.48071> PMID: 26161062
- [14] Lim CHL, Stapleton F, Mehta JS. Review of contact lens-related complications. *Eye Contact Lens* 2018; 44(2): S1-S10. <http://dx.doi.org/10.1097/ICL.0000000000000481> PMID: 29373389
- [15] Konne NM, Collier SA, Spangler J, Cope JR. Healthy contact lens behaviors communicated by eye care providers and recalled by patients — United States, 2018. *MMWR Morb Mortal Wkly Rep* 2019; 68(32): 693-7. <http://dx.doi.org/10.15585/mmwr.mm6832a2> PMID: 31415490
- [16] Liaqat I, Saleem QA, Tahir HM, Arshad M, Arshad N. Identification of virulence factors in contact lens associated bacteria: A physiological approach. *Cont Lens Anterior Eye* 2019; 42(2): 159-64. <http://dx.doi.org/10.1016/j.clae.2018.10.002> PMID: 30337140
- [17] Young G, Veys J, Pritchard N, Coleman S. A multi-centre study of lapsed contact lens wearers. *Ophthalmic Physiol Opt* 2002; 22(6): 516-27. <http://dx.doi.org/10.1046/j.1475-1313.2002.00066.x> PMID: 12477016
- [18] Liesegang TJ. Contact lens-related microbial keratitis: Part I: Epidemiology. *Cornea* 1997; 16(2): 125-31. PMID: 9071523
- [19] Abahussin M, AlAnazi M, Ogbuehi KC, Osuagwu UL. Prevalence, use and sale of contact lenses in Saudi Arabia: Survey on university women and non-ophthalmic stores. *Cont Lens Anterior Eye* 2014; 37(3): 185-90. <http://dx.doi.org/10.1016/j.clae.2013.10.001> PMID: 24211011
- [20] Sulley A, Young G, Hunt C. Factors in the success of new contact lens wearers. *Cont Lens Anterior Eye* 2017; 40(1): 15-24. <http://dx.doi.org/10.1016/j.clae.2016.10.002> PMID: 27818113
- [21] Dumbleton K, Woods CA, Jones LW, Fonn D. The impact of contemporary contact lenses on contact lens discontinuation. *Eye Contact Lens* 2013; 39(1): 93-9. <http://dx.doi.org/10.1097/ICL.0b013e318271caf4> PMID: 23266586
- [22] Richdale K, Sinnott LT, Skadahl E, Nichols JJ. Frequency of and factors associated with contact lens dissatisfaction and discontinuation. *Cornea* 2007; 26(2): 168-74. <http://dx.doi.org/10.1097/01.icc.0000248382.32143.86> PMID: 17251807
- [23] Maruyama K, Yokoi N, Takamata A, Kinoshita S. Effect of environmental conditions on tear dynamics in soft contact lens wearers. *Invest Ophthalmol Vis Sci* 2004; 45(8): 2563-8. <http://dx.doi.org/10.1167/iovs.03-1185> PMID: 15277478
- [24] Chlasta-Twardzik E, Górecka-Nitoń A, Nowińska A, Wylęgała E. The influence of work environment factors on the ocular surface in a one-year follow-up prospective clinical study. *Diagnostics* 2021; 11(3): 392. <http://dx.doi.org/10.3390/diagnostics11030392> PMID: 33668951
- [25] Chaudhary S, Ghimire D, Basu S, Agrawal V, Jacobs DS, Shanbhag SS. Contact lenses in dry eye disease and associated ocular surface disorders. *Indian J Ophthalmol* 2023; 71(4): 1142-53. [http://dx.doi.org/10.4103/IJO.IJO\\_2778\\_22](http://dx.doi.org/10.4103/IJO.IJO_2778_22) PMID: 37026246
- [26] Riley C, Young G, Chalmers R. Prevalence of ocular surface symptoms, signs, and uncomfortable hours of wear in contact lens wearers: The effect of refitting with daily-wear silicone hydrogel lenses (senofilcon a). *Eye Contact Lens* 2006; 32(6): 281-6. <http://dx.doi.org/10.1097/01.icl.0000224522.04723.7a> PMID: 17099389
- [27] Schafer J, Mitchell GL, Chalmers RL, et al. The stability of dryness symptoms after refitting with silicone hydrogel contact lenses over 3 years. *Eye Contact Lens* 2007; 33(5): 247-52. <http://dx.doi.org/10.1097/ICL.0b013e3180587e21> PMID: 17873628
- [28] Moezzi AM, Varikooty J, Luensmann D, et al. The short-term physiological impact of switching reusable silicone hydrogel wearers into a hydrogel daily disposable multifocal. *Clin Ophthalmol* 2019; 13: 1193-202. <http://dx.doi.org/10.2147/OPHTH.S208905> PMID: 31371916
- [29] Bamahfouz AY, Hego NH, Jouhary S, et al. Awareness of contact lens care among college students in Saudi Arabia. *Int J Sci Stud* 2016; 4: 90-6. <http://dx.doi.org/10.17354/ijss/2016/195>
- [30] Boqursain S, Hussain AA, Mubarak AAA, Bujays AD, Mustahi AM. The attitude and awareness of contact lens use among medical students of King Faisal University, Al Ahsa, Saudi Arabia. *J Family Med Prim Care* 2021; 10(10): 3765-71. [http://dx.doi.org/10.4103/jfmpc.jfmpc\\_707\\_21](http://dx.doi.org/10.4103/jfmpc.jfmpc_707_21) PMID: 34934678
- [31] van Hoeven LR, Janssen MP, Roes KCB, Koffijberg H. Aiming for a representative sample: Simulating random versus purposive strategies for hospital selection. *BMC Med Res Methodol* 2015; 15(1): 90. <http://dx.doi.org/10.1186/s12874-015-0089-8> PMID: 26497748
- [32] Astivia OOL, Kroc E, Zumbo BD. The role of item distributions on reliability estimation: The case of cronbach's coefficient alpha. *Educ Psychol Meas* 2020; 80(5): 825-46. <http://dx.doi.org/10.1177/0013164420903770> PMID: 32855561
- [33] Zeried FM, Alnehi DA, Osuagwu UL. A survey on knowledge and attitude of Saudi female students toward refractive correction. *Clin Exp Optom* 2020; 103(2): 184-91. <http://dx.doi.org/10.1111/cxo.12919> PMID: 31115098
- [34] Zhu Q, Yang B, Deng N, et al. The use of contact lenses among university students in Chengdu: Knowledge and practice of contact lens wearers. *Cont Lens Anterior Eye* 2018; 41(2): 229-33. <http://dx.doi.org/10.1016/j.clae.2017.12.008> PMID: 29221709
- [35] Cope JR, Collier SA, Srinivasan K, et al. Contact lens-related corneal infections — United States, 2005–2015. *MMWR Morb Mortal Wkly Rep* 2016; 65(32): 817-20. <http://dx.doi.org/10.15585/mmwr.mm6532a2> PMID: 27538244
- [36] Bakkar MM, Alzghoul EA. Assessment of compliance with contact lens wear and care among university-based population in Jordan.



- Cont Lens Anterior Eye 2020; 43(4): 395-401.  
<http://dx.doi.org/10.1016/j.clae.2020.02.020> PMID: 32127286
- [37] Tajunisah I, Ophth M, Reddy SC, Phuah SJ. Knowledge and practice of contact lens wear and care among medical students of University of Malaya. *Med J Malaysia* 2008; 63(3): 207-10. PMID: 19248691
- [38] Vidotti VG, Kamegasawa A. Profile of medical students from the universidade estadual paulista-UNESP--Botucatu, who wear contact lenses. *Arq Bras Oftalmol* 2006; 69(2): 197-201. <http://dx.doi.org/10.1590/S0004-27492006000200012> PMID: 16699670
- [39] Bhandari M, Hung PR. Habits of contact lens wearers toward lens care in Malaysia. *Med J Malaysia* 2012; 67(3): 274-7. PMID: 23082416
- [40] Beshtawi IM, Qaddumi J, Suboh N, Zaid A, Mansour H, Zeyadeh T. Compliance of soft contact lens care and bacterial contamination among university students in Palestine. *Clin Ophthalmol* 2022; 16: 4121-34. <http://dx.doi.org/10.2147/OPHTH.S352209> PMID: 36536924
- [41] Brant A, Kolomeyer N, Goldberg JL, *et al.* United States population disparities in ophthalmic care. *Ophthalmology* 2023; 130(11): 1121-37. <http://dx.doi.org/10.1016/j.ophtha.2023.06.011> PMID: 37331480
- [42] Sapkota K. Level of compliance in contact lens wearing medical doctors in Nepal. *Cont Lens Anterior Eye* 2015; 38(6): 456-60. <http://dx.doi.org/10.1016/j.clae.2015.05.010> PMID: 26048663
- [43] Lee SY, Kim YH, Johnson D, Mondino BJ, Weissman BA. Contact lens complications in an urgent-care population: The University of California, Los Angeles, contact lens study. *Eye Contact Lens* 2012; 38(1): 49-52. <http://dx.doi.org/10.1097/ICL.0b013e31823ff20e>
- [44] Bergenske P, Long B, Dillehay S, *et al.* Long-term clinical results: 3 years of up to 30-night continuous wear of lotrafilcon A silicone hydrogel and daily wear of low-Dk/t hydrogel lenses. *Eye Contact Lens* 2007; 33(2): 74-80. <http://dx.doi.org/10.1097/01.ICL.0000258591.35468.2c> PMID: 17496699