



The Role of Stress Reduction and Mindfulness Meditation in Glaucoma Management: A Review

Ashley Polski^{1,*} , Anthony P. Mai¹, Brian C. Stagg¹ and Barbara M. Wirostko¹

¹Department of Ophthalmology, Moran Eye Center, University of Utah, Salt Lake City, UT, USA

Abstract:

Glaucoma is a progressive optic neuropathy that is increasing in prevalence worldwide and can lead to devastating vision loss and even blindness if left untreated. Current medical and surgical approaches to glaucoma treatment all aim to lower intraocular pressure (IOP), which is a well-known risk factor for glaucoma development and progression. Although the exact pathogenesis of glaucoma is not yet fully elucidated, multiple studies suggest that mental stress and elevated cortisol levels may contribute to IOP elevation and glaucomatous progression. In recent decades, various randomized-controlled trials have also demonstrated a significant reduction of IOP in patients practicing stress-reduction techniques, including mindfulness meditation and yoga. In this review, we discuss the relationship between mental stress and glaucoma and review the supporting literature to highlight the potential role of mindfulness meditation as an adjunctive strategy to reduce stress and thereby lower IOP in glaucoma patients.

Keywords: Glaucoma, Intraocular pressure, Meditation, Mindfulness, Stress, Yoga.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Department of Ophthalmology, Moran Eye Center, University of Utah, Salt Lake City, UT, USA; E-mail: ashley.polski@hsc.utah.edu

Cite as: Polski A, Mai A, Stagg B, Wirostko B. The Role of Stress Reduction and Mindfulness Meditation in Glaucoma Management: A Review. *Open Ophthalmol J*, 2024; 18: e18743641340238. <http://dx.doi.org/10.2174/0118743641340238241210155401>



Received: June 27, 2024
Revised: November 14, 2024
Accepted: November 19, 2024
Published: December 27, 2024



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Glaucoma is a type of progressive optic neuropathy that affects more than 70 million people, making it the leading cause of irreversible blindness worldwide [1]. Although the underlying pathogenesis of glaucoma remains poorly understood, intraocular pressure (IOP) is a well-established continuous risk factor for glaucomatous damage [2, 3]. Multiple studies, for instance, have demonstrated a direct correlation between increased IOP and the risk of glaucoma development and progression [4-8]. For this reason, medical and surgical glaucoma management, including topical eye drops, laser trabeculoplasty, and incisional glaucoma surgeries, all aim to lower IOP [9, 10].

The prevalence of glaucoma is expected to substantially increase in the coming decades due to an aging population [11]. Consequently, there is growing interest in identifying alternative IOP reduction strategies that could supplement

our current medical and surgical glaucoma treatments [12, 13]. Previously proposed lifestyle-based approaches to IOP reduction include aerobic exercise, stress reduction, and supplementation with antioxidants and neuroprotective agents, such as *Ginkgo biloba*, coenzyme Q10, and nicotinamide [14-16]. In recent years, mindfulness meditation has garnered increased attention as a potential adjunctive treatment for reducing stress and lowering IOP, fueled by emerging evidence linking mental stress to glaucomatous progression [17-20]. However, there is still limited understanding of the effectiveness of mindfulness meditation for IOP reduction and the role it should play in glaucoma management. In this review, we aim to discuss current literature regarding the association between mental stress and glaucoma. We specifically highlight mindfulness meditation and relaxation techniques as a potential complementary approach to lowering IOP in conjunction with standard medical and surgical modalities of glaucoma treatment.

2. RELATIONSHIP BETWEEN MENTAL STRESS AND GLAUCOMA

The pathophysiology of glaucoma is complex and not yet fully elucidated. IOP is a known, modifiable contributor to the development and progression of glaucomatous damage [2, 3]. Vascular dysregulation and autonomic nervous system imbalance have also been suggested as key factors in the pathogenesis of certain glaucoma subtypes, including normal tension glaucoma [21-23]. In recent years, multiple studies have highlighted psychological stress as an upstream factor that can potentially modulate glaucoma risk by increasing IOP and impairing vascular autoregulation [24-29]. Nicou *et al.* (2021) and Turner *et al.* (2019) demonstrated that environmental disturbances causing acute stress led to substantial elevations of IOP in rats and non-human primates, respectively [25, 26]. Similar findings have been demonstrated in human populations. For example, Abe *et al.* (2020) measured significant increases in IOP after inducing psychological stress with the Trier Social Stress Test, a tool used to evaluate stress responses related to public speaking [27]. Jimenez and Vera (2018) similarly showed higher IOP values and cardiovascular indices (such as heart rate and blood pressure) in university students exposed to mental stress induced by test-taking compared to a control group [28]. In a prospective observational study on 151 glaucoma patients, the presence of mental stress was associated not only with increased IOP values but also with worsening of glaucoma based on visual field changes [29]. Such findings have led to an understanding that mental stress itself is a potentially modifiable risk factor for vision loss in glaucoma patients [24].

There are multiple proposed mechanisms for how mental stress modulates IOP and impacts glaucoma pathogenesis based on the physiological adaptations of the body to stress. The typical stress response involves the activation of two neural networks: the sympathetic adrenomedullary (SAM) system and the neuroendocrine stress response system, also known as the hypothalamic-pituitary-adrenal (HPA) axis [24]. The SAM system is responsible for acutely constricting blood vessels and increasing heart rate in response to stress [30]. The HPA axis, on the other hand, produces a slower, more prolonged stress response through the release of glucocorticoids from the adrenal gland into the bloodstream [31]. The activation of these neural networks is thought to influence glaucoma risk by causing intraocular pressure elevation, autonomic nervous system imbalance, and vascular dysregulation [29, 32, 33].

In humans, cortisol is the steroid hormone that is synthesized and released by the adrenal glands in response to HPA axis activation. In addition to regulating the body's stress response, this glucocorticoid hormone plays an important role in suppressing inflammation, regulating blood pressure, modulating the sleep-wake cycle, and regulating the metabolism of nutrients, such as lipids, carbohydrates, and proteins [34]. In the context of chronic psychological stress, elevated levels of cortisol are

associated with a wide variety of systemic conditions, including obesity, cardiovascular disease, diabetes, autoimmune disease, asthma, and cognitive disorders [31, 35, 36]. The effects of cortisol impact the eye as well, with multiple studies dating back over 50 years that have identified a strong correlation between high plasma cortisol levels and the incidence of ocular hypertension and primary open-angle glaucoma [37-40]. Although the anti-inflammatory and metabolic effects of cortisol are crucial in maintaining eye health and have even been harnessed to treat inflammatory eye diseases through the use of topical, periocular, and intravitreal steroid medications [41], prolonged exposure to glucocorticoids can lead to structural and molecular changes in the eye that result in IOP elevation and glaucoma [34]. For example, multiple studies have shown that glucocorticoids contribute to microstructural alterations in the aqueous outflow pathway through the upregulation of myocilin, a protein of unknown function that resides primarily in trabecular meshwork tissue [42, 43]. The binding of cortisol and other steroid medications to glucocorticoid receptor beta has also been postulated to alter trabecular meshwork function through a reduction in phagocytic cell activity [44]. These structural changes associated with cortisol exposure promote aqueous outflow resistance, which leads to elevated IOP and subsequent glaucomatous damage if not controlled appropriately [45, 46].

Even in patients with relatively normal IOP, vascular dysregulation and autonomic imbalance associated with chronic mental stress may play a crucial role in glaucoma pathogenesis, particularly in the setting of normal tension glaucoma [24]. In addition to glucocorticoids, SAM system signaling molecules, such as pro-inflammatory cytokines and endothelin-1, can accelerate the progression of both primary open angle and normal tension glaucoma by contributing to vasoconstriction, endothelial cell dysfunction, and impaired ocular blood flow [24, 47]. Stress hormones may also disrupt nitric oxide signaling, which normally functions as a potent vasodilator and helps to regulate aqueous outflow through the trabecular meshwork [48, 49]. Over time, these vascular changes can result in impaired regulation of ocular perfusion pressure and increased risk of the optic nerve and retinal ganglion cell ischemia, making the eye more vulnerable to glaucomatous injury [32, 33].

The relationship between mental stress and glaucoma is not unidirectional. While chronic psychological stress is linked to an increased risk of glaucoma development and progression, glaucoma itself can exacerbate stress levels. In a study on 589 glaucoma patients, over 80% reported experiencing negative emotions when they received a glaucoma diagnosis [50]. Other studies have demonstrated increased levels of depression and anxiety in patients with progressing or severe glaucoma [51, 52]. Additionally, glaucoma management is complex and often requires regular monitoring, strict medication adherence, and, in some cases, invasive treatments, all of which can further heighten stress levels [53]. Therefore, addressing both the physiological and psychological aspects of glaucoma is

crucial to breaking this negative feedback loop and improving the quality of life for glaucoma patients.

3. MINDFULNESS MEDITATION FOR THE REDUCTION OF IOP

Although mindfulness meditation effectively reduces stress [54], it is not commonly used as an adjunctive glaucoma therapy. According to a 2002 Wills Eye Hospital survey of 1027 patients, only 1.8% of patients had ever utilized mindfulness meditation as a complementary approach to glaucoma treatment [13]. This is perhaps due to relatively scarce scientific evidence to support its usage and a limited understanding of how mindfulness meditation could be effectively applied in real-life clinical practice. Herein, we discuss the randomized-controlled trials that have investigated meditation for glaucoma in recent decades and the impact of mindfulness-based yogic exercises on IOP (Table 1). One of the earliest randomized-controlled trials to investigate the relationship between mindfulness practice and IOP was based in Germany in 1995 [55]. Kaluza and Stempel (1995) randomized primary open-angle glaucoma patients on standard medical therapy to either 1) a training group that underwent eight 90-minute weekly sessions of autogenic relaxation training in addition to their glaucoma drops (n = 11) or 2) a waiting-list control group that remained on their glaucoma drops alone (n = 12). After 8 weeks, both the training group and the control group were exposed to a mental stressor test (MST) designed to increase psychological stress [56]. Although there was no difference in the immediate rise of IOP in response to the MST for patients in the training and control groups ($p > 0.05$), patients in the training group demonstrated an average reduction of IOP by 3 mm Hg compared to the control group ($p < 0.001$) [55, 56]. Unfortunately, the sample size in this study was very small, making it difficult to draw definitive conclusions about the role that mindfulness techniques should play in glaucoma management.

Decades later, Dada *et al.* conducted a series of three larger randomized-controlled trials to investigate the effectiveness of mindfulness meditation in treating glaucoma [18-20]. These studies evaluated the impact of meditation on IOP reduction and also investigated how meditation-induced IOP changes might be explained by changes in serum biomarkers, serum gene expression, trabecular meshwork gene expression, and optic nerve vasculature. In 2018, Dada *et al.* investigated the effect of meditation on primary open-angle glaucoma patients with medication-controlled IOPs [18]. They performed a prospective, single-blinded, and randomized-controlled trial with mindfulness meditation as the intervention arm. The primary endpoint was IOP, while the secondary endpoints were quality of life (QOL) scores (World Health Organization (WHO) questionnaires), stress-related serum biomarkers (*i.e.*, cortisol, β -endorphins, IL6, TNF- α , brain-derived neurotrophic factor (BDNF), reactive oxygen species (ROS), and total antioxidant capacity (TAC)), and whole genome expression. The study included patients

older than 45 years old with IOPs between 12 and 21 mm Hg and a best corrected visual acuity of 20/40 or better. Patients with newly diagnosed or uncontrolled glaucoma, a comorbid condition of vision loss (*i.e.*, macular degeneration), a history of eye surgery in the past 6 months, or prior meditation experience were excluded. Ninety patients were randomized to two groups: 1) meditation + medication (n = 45) and 2) medication only (n = 45). Group 1 did 60 minutes of guided meditation with a trained Mindfulness Stress Based Reduction (MSBR) instructor at 8 AM daily. The meditation sessions consisted of breathing exercises, followed by mindfulness meditation. The study endpoints were measured on the first day of the meditation course and then again on day 21. Of note, the IOP was measured 5 hours after the meditation sessions to avoid possible transient IOP changes directly related to the sessions.

After 3 weeks, the intervention arm had several dramatic results compared to the control group. Analysis revealed a statistically significant reduction in IOP of 6 mm Hg (32%) in the meditation group compared to a reduction of < 1 mm Hg for the non-meditation group ($p < 0.001$). Notably, 30 of the 40 participants (75%) who completed the full meditation course demonstrated a $> 25\%$ reduction in IOP. There was also a significant improvement in the WHO QOL scores in all domains for the intervention arm ($p < 0.001$). Next, blood samples on days 1 and 21 were compared for differences in biochemical markers and gene expression profiling. The meditation group had significant decreases in serum cortisol, IL6, TNF- α , and ROS and significant increases in serum BDNF and TAC. They also had significantly different expressions of 109 genes, of which 54 were upregulated and 55 downregulated. For example, *NGFR*, *TAZ*, *BNP*, *IL2*, *IL4*, *FGFR1*, and metallothionein-I were upregulated, whereas *RAR*, *CYP26A1*, *I-kB*, *EGFR*, *ERK7*, *PTGER3*, and *IL21A* genes were downregulated in the meditation group. The meditation group also had decreased whole genome expression of *BCL2L11*, *CARD8*, and *MAPK10*, all of which are associated with apoptosis and inflammation [18].

This study by Dada *et al.* (2018) not only showed that meditation significantly reduced IOP but also simultaneously improved patient QOL [18]. The IOP reduction in their meditation study arm was comparable to using an additional eye drop ($\sim 30\%$ reduction) without the drop-related side effects or QOL reduction [57]. The improved QOL parameters in the meditation group suggest that meditation may effectively target the mental stress that is thought to be an upstream factor for glaucoma risk. Dada *et al.* (2018) then used serum biomarkers and whole genome expression to build a possible mechanistic framework for how meditation ultimately impacts IOP [18]. For instance, the meditation group had decreased serum cortisol, IL6, TNF- α , and ROS levels. Cortisol, like other glucocorticoids, is notorious for increasing IOP by altering trabecular meshwork microstructure, as previously discussed. IL6 and TNF- α are both inflammatory markers, the latter of which causes

metalloproteinase-associated tissue remodeling in glaucomatous optic nerve heads and IOP-induced retinal ganglion cell death [58, 59]. Additionally, oxidative stress from ROS damages the trabecular meshwork, especially the endothelial cells, which can further raise IOP [60, 61]. Therefore, a reduction in these biomarkers after mindfulness meditation may have contributed to the lower IOP seen in the intervention arm. In summary, meditation appeared to reduce stress-related serum biomarkers and modulate the expression of genes that may have harmful effects on the trabecular meshwork through inflammation, tissue remodeling, and apoptosis. The conclusion that these systemic changes affected ocular tissues in this study was an assumption until the second study in 2021 addressed it [20].

In 2021, Dada *et al.* subsequently conducted a randomized-controlled trial that assessed primary open-angle glaucoma patients in imminent need of a trabeculectomy due to uncontrolled high IOP [20]. This study expanded on the conclusions of the 2018 study by

also exploring if meditation actually affected ocular tissues. The 2021 study included patients with IOPs ≥ 21 mm Hg and optic nerve head cup-to-disc ratio >0.7 on maximum medical therapy. The endpoints were similar to the 2018 study but also included diurnal IOP measurements (Goldmann tonometry every 3 hours on days 1 and 21) [18, 20]. The 2021 study was conducted in two stages (Fig. 1). In stage 1, 60 patients were randomized to an intervention group (daily guided meditation + current medications) and a control group (current medications only) for 21 days. On day 21, “success” was defined as IOP ≤ 15 mm Hg and “failure” as IOP >15 mm Hg. In stage 2, those with IOP ≤ 15 mm Hg continued their current management, while those who failed received a trabeculectomy and had their trabecular meshwork punches submitted for genome expression profiling. The patients who initially were in the meditation arm, regardless of whether trabeculectomy was ultimately performed, were instructed to continue meditating at home without guidance. Repeat measurements were done after 6 weeks (9 weeks total after enrollment).

Table 1. Summary of the participants, interventions, and main findings of randomized-controlled trials that have studied the role of meditation and relaxation techniques in intraocular pressure (IOP) reduction for Primary Open-Angle Glaucoma (POAG) patients.

Authors/Year	Participants and Intervention	Main Results
Kaluza and Stempel 1995-6 [55, 56]	23 POAG patients on standard drops: <ul style="list-style-type: none"> • 11 patients in the study group underwent 90-minute weekly sessions of autogenic relaxation training for 8 weeks in addition to drops • 12 patients in the control group used drops alone 	Average reduction of IOP by 3 mm Hg in the study group compared to the control group ($p < 0.001$)
Gagrani <i>et al.</i> 2018 [71]	60 POAG patients on standard drops: <ul style="list-style-type: none"> • 30 patients in the study group underwent 45-minute daily breathing exercises for 6 weeks in addition to drops • 30 patients in the control group used drops alone 	Average reduction of IOP by 1.5 mm Hg in the study group compared to <0.1 mm Hg in the control group ($p = 0001$)
Dada <i>et al.</i> 2018 [18]	90 POAG patients on standard drops: <ul style="list-style-type: none"> • 45 patients in the study group underwent 1 hour of daily mindfulness meditation for 3 weeks in addition to drops • 45 patients in the control group used drops alone 	Average reduction of IOP by 6 mm Hg in the study group compared to <1 mm Hg in the control group ($p < 0.001$)
Dada <i>et al.</i> 2021 [20]	60 POAG patients with uncontrolled high IOP on maximum drops: <p><u>Stage 1:</u></p> <ul style="list-style-type: none"> • 30 patients in the study group underwent daily guided meditation for 3 weeks in addition to drops • 30 patients in the control group used drops alone <p><u>Stage 2:</u></p> <ul style="list-style-type: none"> • Patients with IOP ≤ 15 mm Hg continued current management • Patients with IOP >15 mm Hg underwent trabeculectomy in addition to home meditation or drops 	<p><u>Stage 1:</u></p> <p>Average reduction of IOP by 5 mm Hg in the study group compared to 0.2 mm Hg in the control group ($p = 0.001$)</p> <p><u>Stage 2:</u></p> <p>50% of participants in the study (meditation) group had IOP >15 mm Hg and required trabeculectomy, and 100% of participants in the control (non-meditation) group had IOP >15 mm Hg and required trabeculectomy</p>
Dada <i>et al.</i> 2021 [19]	60 POAG patients on standard drops: <ul style="list-style-type: none"> • 30 patients in the study group underwent a 6-week mindfulness meditation course in addition to drops • 30 patients in the control group used drops alone • Circumpapillary vessel density was evaluated with optical coherence tomography angiography (OCT-A) in both groups 	Average reduction of IOP by 2 mm Hg in the study group compared to 0.8 mm Hg in the control group ($p = 0.001$) and increased circumpapillary vessel density on OCT-A in the study group compared to the control group
Udenia <i>et al.</i> 2021 [69]	90 POAG patients on standard drops: <ul style="list-style-type: none"> • 45 patients in the study group underwent 30-minute daily breathing exercises in addition to drops • 45 patients in the control group used drops alone 	Average reduction of IOP by 6 mm Hg in the study group compared to the control group ($p < 0.001$)
Ismail <i>et al.</i> 2022 [70]	62 POAG patients on standard drops: <ul style="list-style-type: none"> • 31 patients in the study group underwent daily guided <i>Jyoti-Trataka</i> mindfulness sessions in addition to drops • 31 patients in the control group used drops alone 	Average reduction of IOP by 3 mm Hg in the study group and 0 mm Hg in the control group ($p < 0.001$)




Fig. (1). Flow chart of the study by Dada et al. (2021).

After 21 days and at the end of stage 1, the meditation group had a significant decrease in mean IOP (5 mm Hg, 23.34% reduction) and diurnal IOP range (4.07 to 1.75 mm Hg) compared to that of the control group (mean IOP decrease of 0.2 mmHg and change in diurnal IOP range from 4.5 to 4.28 mm Hg; $p = 0.001$). Half of the participants in the meditation group ($n = 15$) and all of the participants in the control group ($n = 30$) did not show any significant change ($IOP > 15$ mm Hg). Of note, 7 patients in the meditation group who met the success criteria ($IOP \leq 15$ mm Hg) actually had $IOP \leq 12$ mm Hg at day 21. At the end of stage 2, there were patients in 3 groups: meditation + drops, meditation + trabeculectomy, and trabeculectomy only (Fig. 1). All groups had significant reductions in IOP by the end of the study period (mean IOP of 12.8 mmHg, 11.8 mmHg, and 13.3 mmHg, respectively; $p = 0.0001$). The patients with the largest ultimate IOP decrease were those in the meditation group who also received a trabeculectomy. Trabecular meshwork gene expression was compared between the intervention and control groups; the meditators had significant upregulation of *NGB*, *NGFR*, *NRG1*, *NOS1*, *NOS3*, *MNTR1A*, *NR3C1*, and *MT1E*, and downregulation of *IL4*, *BCL2L11*, *CARD8*, *FGFR1*, *MAPK10*, *MAPK15*, *TGFB*, *TNFA*, *NFkB*, and *EGFR*.

This study showed that meditation was associated with both significantly decreased mean IOP and diurnal IOP range [20]. About 86% of the meditation group had $\geq 15\%$ IOP reduction, and half of the meditation group had enough IOP reduction to avoid trabeculectomy. This effect was sustained 6 weeks later, even though the guided group meditation session was changed to home meditation, implying that patients were perhaps able to maintain an effective meditation practice by themselves.

Even the meditators who needed a trabeculectomy had the largest IOP decrease compared to the patients with trabeculectomy alone. These results suggest that meditation may also be effective as an adjunctive therapy for patients with uncontrolled IOP; not only can it potentially delay incisional surgery, but it can also be a supplementary treatment pre- and post-operatively if surgery is indicated. Lastly, the comparison of the trabecular meshwork genome expression revealed that gene expression in ocular tissues between the intervention and control groups had significant differences. The meditation group had decreased expression of *NFkB* (pro-inflammation), *TGFB* (trabecular meshwork fibrosis), and *EGFR* (trabecular meshwork remodeling) and increased expression of *NOS* (increased trabecular meshwork flow and vasodilation) [48] and melatonin receptors (IOP homeostasis) [20]. This 2021 study strengthens the hypothesis that meditation first reduces the whole body stress response (serum biomarker and whole genome expression) [18], which then changes ocular tissues (trabecular meshwork genome expression) to eventually reduce IOP [20].

In 2021, Dada et al. also performed a follow-up randomized controlled trial investigating the effects of meditation on the optic nerve head using ocular coherence tomography angiography (OCT-A) [19]. As vascular changes have been linked to glaucoma [22, 62], this study evaluated the effect of meditation on the optic nerve head microvasculature. The study design was similar to that of a study conducted in 2018; 60 primary open-angle glaucoma patients with IOPs < 21 mm Hg on medications were randomized to meditation and control groups [18, 19]. OCT-A and IOP measurements were performed at the beginning and end of a 6-week meditation course. The

study again confirmed that daily meditation significantly reduced IOP compared to the control group ($p = 0.001$), with 23 (38%) eyes demonstrating a <10% reduction in IOP, 33 (55%) eyes showing a 10-15% reduction in IOP, and 4 (7%) eyes having a >15% reduction in IOP. OCT-A imaging showed that the meditation group had significantly increased circumpapillary vessel density superiorly ($p = 0.02$) and nasally ($p = 0.01$), circumpapillary vessel perfusion ($p < 0.001$), and flux index ($p < 0.001$) compared to the control group. As decreased circumpapillary vessel perfusion and density have been linked to worsening visual fields, increasing these parameters via meditation may be protective against glaucoma progression [63, 64]. Although the mechanism for how meditation specifically affects optic nerve head vasculature has not yet been elucidated, it is reasonable to postulate that meditation-induced reduction of SAM system signaling, as previously discussed, could counteract sympathetic-driven vasoconstriction and lead to increased vascular perfusion of the nerve [24, 47].

Another practice that has been studied in relation to glaucoma and IOP reduction is yoga, which incorporates physical postures, breathing techniques, focusing exercises, and mindfulness meditation to reduce stress and foster physical and mental well-being [65-68]. Yogic breathing and focusing exercises, such as *Jyoti-Trataka* (gazing at a single point or candle flame) or meditation-focused breathing, have been shown to reduce IOP in glaucoma patients [69-71]. In a clinical trial by Ismail *et al.* (2022), 62 open-angle glaucoma patients were included. Thirty-one patients were randomized to a study group that participated in guided *Jyoti-Trataka* mindfulness sessions with a certified yoga instructor in addition to their baseline glaucoma medication, and 31 patients were randomized to a control group with glaucoma medication only [70]. Compared to the control group, which demonstrated a mean IOP of 26.40 mm Hg after 1 month, the study arm demonstrated a significant reduction in not only mean IOP (23.23 mm Hg) but also systolic and diastolic blood pressure measurements ($p < 0.05$). This study supported the hypothesis that yogic focusing exercises, such as *Jyoti-Trataka*, may be useful in reducing IOP in glaucoma patients in conjunction with standard medical therapy for glaucoma. Yogic breathing exercises have also been associated with a potentially beneficial effect on IOP in glaucoma patients. In a randomized-controlled trial by Udenia *et al.* (2021), 90 patients with moderate and severe primary open-angle glaucoma were randomized to either a yoga group combined with glaucoma medication ($n = 45$) or a control group with glaucoma medication only ($n = 45$) [69]. Patients in the yoga group participated in diaphragmatic breathing and alternate nostril breathing exercises for 30 minutes per day over a period of 6 months. Compared to the control group, the yoga group demonstrated IOP reductions of nearly 6 mm Hg in each eye after 6 months ($p < 0.001$). Gagrani *et al.* (2018) also investigated the effect of breathing exercises on IOP in moderate and severe primary open-angle glaucoma patients [71]. In their study

on 60 glaucoma patients, 30 patients were randomized to a yoga group in addition to standard medical treatment, and 30 patients were randomized to a control group with standard medical treatment only. The yoga group participated in meditation focused on breathing exercises for 45 minutes per day over 6 weeks. After 6 weeks, the yoga group demonstrated a significant reduction in mean IOP (15.9 to 14.4 mm Hg; $p = 0.0001$) compared to the control group (15.7 to 15.65 mm Hg; $p = 0.41$) [71]. These studies demonstrated that mindfulness-based breathing exercises may be useful for reducing IOP in both the short-term (6 weeks) [71] and the long-term (6 months) [69].

Unfortunately, not all relaxation practices are helpful for achieving IOP reduction, and some forms of yoga may even lead to elevated IOP and increased risk of glaucoma progression [65]. For example, two case studies and a prospective single-center observational study showed that breath-holding exercises and *Asanas* (yoga postures) that involve head-down positioning are associated with significantly increased IOP during practice and, in some cases, may even lead to worsening visual field defects [72-74]. For this reason, providers should be prudent in the types of yoga and mindfulness practices that they recommend to patients with glaucoma.

4. LIMITATIONS AND FUTURE DIRECTIONS

The studies discussed in this review highlight the possible effectiveness of mindfulness meditation and relaxation techniques as adjunctive strategies for IOP reduction. However, there is still a scarcity of data regarding the usage of meditation in actual clinical practice, as well as a limited understanding of how these techniques impact IOP and glaucoma prognosis in the long term. Many of the studies that have investigated the relationship between mindfulness meditation, yogic practices, and IOP reduction have been limited to relatively small sample sizes. For example, Kaluza and Stempel (1995) included only 23 patients in total, which limits the generalizability and reliability of their findings [55]. To address this sample size limitation and provide a more precise estimate of the relationship between mindfulness practices and IOP, two meta-analyses were performed. Zaher *et al.* (2023) evaluated studies related to the impact of meditation, autogenic or ocular relaxation exercises, or yoga on IOP, and Chetry *et al.* (2023) evaluated studies related to the impact of yogic practices on IOP [65, 75]. In the meta-analysis by Zaher *et al.* (2023), regular practice of yoga for 6 weeks was associated with a 22% reduction in IOP, and meditation was associated with an average reduction of IOP by 2.02 mm Hg [75]. Similarly, in the meta-analysis by Chetry *et al.* (2023), pooled data from multiple randomized-controlled trials showed a decrease in mean IOP by 1-2 mm Hg after 4-6 weeks of participation in relaxation yoga techniques ($p < 0.05$) [65]. Although meta-analyses can be useful for increasing overall sample size and, consequently, increasing power to study an effect of interest [76], additional primary studies with larger sample sizes would be useful to better understand the

impact of mindfulness practices on IOP in glaucoma patients.

Randomization of patients into exposure and control groups is useful for reducing the potential for bias in a study. However, in the context of mindfulness meditation and IOP-related studies, it is difficult to account for all potential confounding variables since IOP can be influenced by numerous factors. In multiple studies, for example, patients were continued on their baseline topical glaucoma medications, which could potentially confound the IOP findings (although Dada *et al.* attempted to account for this by demonstrating no significant differences between the intervention and control groups with regard to the types ($p = 0.21$ to 0.34) and numbers ($p = 0.19$ to 0.70) of glaucoma medications used) [19, 20, 55, 56, 69, 70]. Additionally, these studies had relatively short follow-up intervals (ranging from 3 to 9 weeks), so it is uncertain if the beneficial effects of meditation could be sustained for months to years [65, 75]. The current lack of long-term follow-up precludes the ability to determine if meditation actually reduces visual field loss, which is the end-point of greatest clinical utility. Therefore, we cannot yet know how the use of mindfulness meditation in clinical practice would impact the long-term visual prognosis of glaucoma patients. Further clinical trials with longer follow-up periods are needed to assess if meditation actually decreases glaucoma progression long-term.

Although there were no harmful effects from mindfulness meditation alone, it was still time-consuming to commit to one hour of meditation daily. For this reason, many studies used an instructor to provide daily guided meditations and ensure consistent compliance with treatment sessions [18-20, 56]. It is unclear if patients have the time or motivation in real-life settings to attend guided hour-long meditation sessions or if this duration of meditation is even necessary to achieve useful IOP reduction. The 2021 study by Dada *et al.* showed encouraging data that after 3 weeks of guided meditation sessions, those who continued their practice at home without guidance maintained a lower IOP even 6 weeks later [20]. A survey of 100 patients in Australia also showed that 63% were amenable to doing 45-60 minutes of meditation daily for glaucoma if prescribed by a doctor [17]. Still, agreeability and actual performance are two different aspects. Further studies that mimic real-world settings, such as fewer guided sessions, shorter meditation times, or even the usage of a meditation-guided phone application, are needed to evaluate the practicality of meditation as a therapy.

Although mindfulness meditation has gained increasing acceptance in the medical community, especially *via* MSBR therapy, it is still not yet considered a mainstream adjunctive medical treatment. There may be a scarcity of trained MSBR instructors or a lack of awareness on how patients can be referred to them. We hope that these logistical hurdles become less as the evidence for meditation as a complementary glaucoma therapy increases.

CONCLUSION

In numerous randomized-controlled trials, mindfulness meditation and relaxation techniques have been associated with a significant reduction of IOP relative to standard medical treatment of glaucoma alone. These findings suggest that meditation may be a useful adjunctive, non-invasive strategy to optimize IOP management in glaucoma patients. However, it is still unclear whether the beneficial effects of meditation persist long-term. Additionally, mindfulness-based practices should not be used as a replacement for standard medical and surgical treatment of glaucoma. Furthermore, multi-year studies are needed to ascertain the real-life clinical feasibility and long-term effectiveness of mindfulness meditation in glaucoma management.

AUTHORS' CONTRIBUTION

A.P. and A.P.M.: Study conception and design; A.P. and A.P.M.: Data collection; A.P. and A.P.M.: Analysis and interpretation of results; A.P., A.P.M., B.C.S., and B.M.W.: Draft manuscript. All authors reviewed the results and approved the final version of the manuscript.

CONSENT FOR PUBLICATION

Not applicable.

FUNDING

This work was financially supported by the Achievement Reward for College Scientists (ARCS) Foundation (AP) and an unrestricted grant from Research to Prevent Blindness (BCS, BMW), which had no role in the writing, editing, approval, or decision to publish this article.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Weinreb RN, Aung T, Medeiros FA. The pathophysiology and treatment of glaucoma: A review. *JAMA* 2014; 311(18): 1901-11. <http://dx.doi.org/10.1001/jama.2014.3192> PMID: 24825645
- [2] Nickells RW, Howell GR, Soto I, John SWM. Under pressure: Cellular and molecular responses during glaucoma, a common neurodegeneration with axonopathy. *Annu Rev Neurosci* 2012; 35(1): 153-79. <http://dx.doi.org/10.1146/annurev.neuro.051508.135728> PMID: 22524788
- [3] Gordon MO, Beiser JA, Brandt JD, *et al.* The ocular hypertension treatment study: Baseline factors that predict the onset of primary open-angle glaucoma. *Arch Ophthalmol* 2002; 120(6): 714-20. <http://dx.doi.org/10.1001/archoph.120.6.714> PMID: 12049575
- [4] Quigley HA. Glaucoma. *Lancet* 2011; 377(9774): 1367-77. [http://dx.doi.org/10.1016/S0140-6736\(10\)61423-7](http://dx.doi.org/10.1016/S0140-6736(10)61423-7) PMID: 21453963
- [5] Wilson M, Singh K. Intraocular pressure: Does it measure up? *Open Ophthalmol J* 2009; 3(1): 32-7. <http://dx.doi.org/10.2174/1874364100903010032> PMID: 19812717

- [6] Sihota R, Angmo D, Ramaswamy D, Dada T. Simplifying “target” intraocular pressure for different stages of primary open-angle glaucoma and primary angle-closure glaucoma. *Indian J Ophthalmol* 2018; 66(4): 495-505. http://dx.doi.org/10.4103/ijo.IJO_1130_17 PMID: 29582808
- [7] Sommer A, Tielsch JM, Katz J, *et al.* Relationship between intraocular pressure and primary open angle glaucoma among white and black Americans. The Baltimore Eye Survey. *Arch Ophthalmol* 1991; 109(8): 1090-5. <http://dx.doi.org/10.1001/archophth.1991.01080080050026> PMID: 1867550
- [8] Varma R, Ying-Lai M, Francis BA, *et al.* Prevalence of open-angle glaucoma and ocular hypertension in latinos. *Ophthalmology* 2004; 111(8): 1439-48. <http://dx.doi.org/10.1016/j.ophtha.2004.01.025> PMID: 15288969
- [9] Heijl A, Leske MC, Bengtsson B, Hyman L, Bengtsson B, Hussein M. Reduction of intraocular pressure and glaucoma progression: Results from the early manifest glaucoma trial. *Arch Ophthalmol* 2002; 120(10): 1268-79. <http://dx.doi.org/10.1001/archophth.120.10.1268> PMID: 12365904
- [10] Kerr NM. The changing glaucoma treatment paradigm. *Clin Exp Ophthalmol* 2022; 50(2): 126-7. <http://dx.doi.org/10.1111/ceo.14052> PMID: 35174602
- [11] Allison K, Patel D, Alabi O. Epidemiology of glaucoma: The past, present, and predictions for the future. *Cureus* 2020; 12(11): e11686. <http://dx.doi.org/10.7759/cureus.11686> PMID: 33391921
- [12] Rhee DJ, Katz LJ, Spaeth GL, Myers JS. Complementary and alternative medicine for glaucoma. *Surv Ophthalmol* 2001; 46(1): 43-55. [http://dx.doi.org/10.1016/S0039-6257\(01\)00233-8](http://dx.doi.org/10.1016/S0039-6257(01)00233-8) PMID: 11525790
- [13] Rhee DJ, Spaeth GL, Myers JS, *et al.* Prevalence of the use of complementary and alternative medicine for glaucoma. *Ophthalmology* 2002; 109(3): 438-43. [http://dx.doi.org/10.1016/S0161-6420\(01\)01030-2](http://dx.doi.org/10.1016/S0161-6420(01)01030-2) PMID: 11874744
- [14] Sim RH, Sirasanagandla SR, Das S, Teoh SL. Treatment of glaucoma with natural products and their mechanism of action: An update. *Nutrients* 2022; 14(3): 534. <http://dx.doi.org/10.3390/nu14030534> PMID: 35276895
- [15] Parikh RS, Parikh SR. Alternative therapy in glaucoma management: Is there any role? *Indian J Ophthalmol* 2011; 59(S1): S158-60. <http://dx.doi.org/10.4103/0301-4738.73679>
- [16] Goulart Nacácio e Silva S, Occhiutto ML, Costa VP, Costa VP. The use of Nicotinamide and Nicotinamide riboside as an adjunct therapy in the treatment of glaucoma. *Eur J Ophthalmol* 2023; 33(5): 1801-15. <http://dx.doi.org/10.1177/11206721231161101> PMID: 36916064
- [17] Brogan K, Bigirimana D, Wightman A, Green C, Martin KR. Daily meditation practice for managing glaucoma patients’ attitudes and acceptance. *J Glaucoma* 2022; 31(9): e75-82. <http://dx.doi.org/10.1097/IJG.0000000000002076> PMID: 35882038
- [18] Dada T, Mittal D, Mohanty K, *et al.* Mindfulness meditation reduces intraocular pressure, lowers stress biomarkers and modulates gene expression in glaucoma: A randomized controlled trial. *J Glaucoma* 2018; 27(12): 1061-7. <http://dx.doi.org/10.1097/IJG.0000000000001088> PMID: 30256277
- [19] Dada T, Lahri B, Mahalingam K, *et al.* Beneficial effect of mindfulness based stress reduction on optic disc perfusion in primary open angle glaucoma: A randomized controlled trial. *J Tradit Complement Med* 2021; 11(6): 581-6. <http://dx.doi.org/10.1016/j.jtcm.2021.06.006> PMID: 34765522
- [20] Dada T, Bhai N, Midha N, *et al.* Effect of mindfulness meditation on intraocular pressure and trabecular meshwork gene expression: A randomized controlled trial. *Am J Ophthalmol* 2021; 223: 308-21. <http://dx.doi.org/10.1016/j.ajo.2020.10.012> PMID: 33393484
- [21] Konieczka K, Choi HJ, Koch S, Fankhauser F, Schoetzau A, Kim DM. Relationship between normal tension glaucoma and Flammer syndrome. *EPMA J* 2017; 8(2): 111-7. <http://dx.doi.org/10.1007/s13167-017-0097-3> PMID: 28725291
- [22] Flammer J, Haefliger IO, Orgül S, Resink T. Vascular dysregulation: A principal risk factor for glaucomatous damage? *J Glaucoma* 1999; 8(3): 212-9. <http://dx.doi.org/10.1097/00061198-199906000-00012> PMID: 10376264
- [23] Grieshaber MC, Mozaffarieh M, Flammer J. What is the link between vascular dysregulation and glaucoma? *Surv Ophthalmol* 2007; 52(6) (Suppl. 2): S144-54. <http://dx.doi.org/10.1016/j.survophthal.2007.08.010> PMID: 17998040
- [24] Sabel BA, Wang J, Cárdenas-Morales L, Faiq M, Heim C. Mental stress as consequence and cause of vision loss: the dawn of psychosomatic ophthalmology for preventive and personalized medicine. *EPMA J* 2018; 9(2): 133-60. <http://dx.doi.org/10.1007/s13167-018-0136-8> PMID: 29896314
- [25] Nicou CM, Pillai A, Passaglia CL. Effects of acute stress, general anesthetics, tonometry, and temperature on intraocular pressure in rats. *Exp Eye Res* 2021; 210: 108727. <http://dx.doi.org/10.1016/j.exer.2021.108727> PMID: 34390732
- [26] Turner DC, Miranda M, Morris JS, Girkin CA, Downs JC. Acute stress increases intraocular pressure in nonhuman primates. *Ophthalmol Glaucoma* 2019; 2(4): 210-4. <http://dx.doi.org/10.1016/j.ogla.2019.03.010> PMID: 31799505
- [27] Abe RY, Silva TC, Dantas I, *et al.* Can psychologic stress elevate intraocular pressure in healthy individuals? *Ophthalmol Glaucoma* 2020; 3(6): 426-33. <http://dx.doi.org/10.1016/j.ogla.2020.06.011> PMID: 32768362
- [28] Jiménez R, Vera J. Effect of examination stress on intraocular pressure in university students. *Appl Ergon* 2018; 67: 252-8. <http://dx.doi.org/10.1016/j.apergo.2017.10.010> PMID: 29122197
- [29] Marc A, Stan C. Effect of physical and psychological stress on the course of primary open angle glaucoma. *Oftalmologia* 2013; 57(2): 60-6. PMID: 24386794
- [30] Ulrich-Lai YM, Herman JP. Neural regulation of endocrine and autonomic stress responses. *Nat Rev Neurosci* 2009; 10(6): 397-409. <http://dx.doi.org/10.1038/nrn2647> PMID: 19469025
- [31] Cohen S, Janicki-Deverts D, Doyle WJ, *et al.* Chronic stress, glucocorticoid receptor resistance, inflammation, and disease risk. *Proc Natl Acad Sci USA* 2012; 109(16): 5995-9. <http://dx.doi.org/10.1073/pnas.1118355109> PMID: 22474371
- [32] Riccadonna M, Covi G, Pancera P, *et al.* Autonomic system activity and 24-hour blood pressure variations in subjects with normal- and high-tension glaucoma. *J Glaucoma* 2003; 12(2): 156-63. <http://dx.doi.org/10.1097/00061198-200304000-00011> PMID: 12671471
- [33] Na KS, Lee NY, Park SH, Park CK. Autonomic dysfunction in normal tension glaucoma: the short-term heart rate variability analysis. *J Glaucoma* 2010; 19(6): 377-81. <http://dx.doi.org/10.1097/IJG.0b013e3181c4ae58> PMID: 20051893
- [34] Patel PD, Kodati B, Clark AF. Role of glucocorticoids and glucocorticoid receptors in glaucoma pathogenesis. *Cells* 2023; 12(20): 2452. <http://dx.doi.org/10.3390/cells12202452> PMID: 37887296
- [35] Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci* 2009; 10(6): 434-45. <http://dx.doi.org/10.1038/nrn2639> PMID: 19401723
- [36] Heim C, Ehlert U, Hellhammer DH. The potential role of hypocortisolism in the pathophysiology of stress-related bodily disorders. *Psychoneuroendocrinology* 2000; 25(1): 1-35. [http://dx.doi.org/10.1016/S0306-4530\(99\)00035-9](http://dx.doi.org/10.1016/S0306-4530(99)00035-9) PMID: 10633533
- [37] Schwartz B, McCarty G, Rosner B. Increased plasma free cortisol

- in ocular hypertension and open angle glaucoma. *Arch Ophthalmol* 1987; 105(8): 1060-5.
<http://dx.doi.org/10.1001/archoph.1987.01060080062029> PMID: 2888454
- [38] Rozsival P, Hampl R, Obenberger J, Stárka L, Řehák S. Aqueous humour and plasma cortisol levels in glaucoma and cataract patients. *Curr Eye Res* 1981; 1(7): 391-6.
<http://dx.doi.org/10.3109/02713688109019976> PMID: 7318492
- [39] Ray S, Mehra KS, Misra S, Singh R. Plasma cortisol in glaucoma. *Ann Ophthalmol* 1977; 9(9): 1151-4.
 PMID: 561563
- [40] Weitzman ED, Henkind P, Leitman M, Hellman L. Correlative 24-hour relationships between intraocular pressure and plasma cortisol in normal subjects and patients with glaucoma. *Br J Ophthalmol* 1975; 59(10): 566-72.
<http://dx.doi.org/10.1136/bjo.59.10.566> PMID: 1191614
- [41] Fung AT, Tran T, Lim LL, *et al.* Local delivery of corticosteroids in clinical ophthalmology: A review. *Clin Exp Ophthalmol* 2020; 48(3): 366-401.
<http://dx.doi.org/10.1111/ceo.13702> PMID: 31860766
- [42] Sharma R, Grover A. Myocilin-associated glaucoma: A historical perspective and recent research progress. *Mol Vis* 2021; 27: 480-93.
 PMID: 34497454
- [43] Resch ZT, Fautsch MP. Glaucoma-associated myocilin: A better understanding but much more to learn. *Exp Eye Res* 2009; 88(4): 704-12.
<http://dx.doi.org/10.1016/j.exer.2008.08.011> PMID: 18804106
- [44] Zhang X, Ognibene CM, Clark AF, Yorio T. Dexamethasone inhibition of trabecular meshwork cell phagocytosis and its modulation by glucocorticoid receptor β . *Exp Eye Res* 2007; 84(2): 275-84.
<http://dx.doi.org/10.1016/j.exer.2006.09.022> PMID: 17126833
- [45] Wei M, Huang ZY, Zhang GW, Guan HJ, Ji M, Ji M. Expression profile analysis to identify potential gene changes induced by dexamethasone in the trabecular meshwork. *Int J Ophthalmol* 2022; 15(8): 1240-8.
<http://dx.doi.org/10.18240/ijo.2022.08.03> PMID: 36017046
- [46] Mohd Nasir NA, Agarwal R, Krasilnikova A, Sheikh Abdul Kadir SH, Iezhitsa I. Effect of dexamethasone on the expression of MMPs, adenosine A1 receptors and NFKB by human trabecular meshwork cells. *J Basic Clin Physiol Pharmacol* 2020; 31(6): 20190373.
<http://dx.doi.org/10.1515/jbcpp-2019-0373> PMID: 32697755
- [47] Toda N, Nakanishi-Toda M. How mental stress affects endothelial function. *Pflugers Arch* 2011; 462(6): 779-94.
<http://dx.doi.org/10.1007/s00424-011-1022-6> PMID: 21947555
- [48] Aliancy J, Stamer WD, Wirostko B. A review of nitric oxide for the treatment of glaucomatous disease. *Ophthalmol Ther* 2017; 6(2): 221-32.
<http://dx.doi.org/10.1007/s40123-017-0094-6> PMID: 28584936
- [49] Schmetterer L, Polak K. Role of nitric oxide in the control of ocular blood flow. *Prog Retin Eye Res* 2001; 20(6): 823-47.
[http://dx.doi.org/10.1016/S1350-9462\(01\)00014-3](http://dx.doi.org/10.1016/S1350-9462(01)00014-3) PMID: 11587919
- [50] Odberg T, Jakobsen JE, Hultgren SJ, Halseide R. The impact of glaucoma on the quality of life of patients in Norway. *Acta Ophthalmol Scand* 2001; 79(2): 116-20.
<http://dx.doi.org/10.1034/j.1600-0420.2001.079002116.x> PMID: 11284746
- [51] Mabuchi F, Yoshimura K, Kashiwagi K, *et al.* Risk factors for anxiety and depression in patients with glaucoma. *Br J Ophthalmol* 2012; 96(6): 821-5.
<http://dx.doi.org/10.1136/bjophthalmol-2011-300910> PMID: 22353697
- [52] Diniz-Filho A, Abe RY, Cho HJ, Baig S, Gracitelli CPB, Medeiros FA. Fast visual field progression is associated with depressive symptoms in patients with glaucoma. *Ophthalmology* 2016; 123(4): 754-9.
<http://dx.doi.org/10.1016/j.ophtha.2015.12.014> PMID: 26920097
- [53] Latif K, Nishida T, Moghimi S, Weinreb RN. Quality of life in glaucoma. *Graefes Arch Clin Exp Ophthalmol* 2023; 261(10): 3023-30.
<http://dx.doi.org/10.1007/s00417-023-06050-z> PMID: 37017741
- [54] Brown KW, Ryan RM. The benefits of being present: Mindfulness and its role in psychological well-being. *J Pers Soc Psychol* 2003; 84(4): 822-48.
<http://dx.doi.org/10.1037/0022-3514.84.4.822> PMID: 12703651
- [55] Kaluza G, Strempele U. Effects of self-relaxation methods and visual imagery on IOP in patients with open-angle glaucoma. *Ophthalmologica* 1995; 209(3): 122-8.
<http://dx.doi.org/10.1159/000310596> PMID: 7630617
- [56] Kaluza G, Strempele I, Maurer H. Stress reactivity of intraocular pressure after relaxation training in open-angle glaucoma patients. *J Behav Med* 1996; 19(6): 587-97.
<http://dx.doi.org/10.1007/BF01904906> PMID: 8970917
- [57] Balkrishnan R, Brent Bond J, Byerly WG, Camacho FT, Anderson RT. Medication-related predictors of health-related quality of life in glaucoma patients enrolled in a medicare health maintenance organization. *Am J Geriatr Pharmacother* 2003; 1(2): 75-81.
[http://dx.doi.org/10.1016/S1543-5946\(03\)90003-1](http://dx.doi.org/10.1016/S1543-5946(03)90003-1) PMID: 15555469
- [58] Akhter N, Nix M, Abdul Y, Singh S, Husain S. Delta-opioid receptors attenuate TNF- α -induced MMP-2 secretion from human ONH astrocytes. *Invest Ophthalmol Vis Sci* 2013; 54(10): 6605-11.
<http://dx.doi.org/10.1167/iov.13-12196> PMID: 24030463
- [59] Zalewska R, Reszeć J, Kisielewski W, Mariak Z. Metalloproteinase 9 and TIMP-1 expression in retina and optic nerve in absolute angle closure glaucoma. *Adv Med Sci* 2016; 61(1): 6-10.
<http://dx.doi.org/10.1016/j.advms.2015.07.007> PMID: 26342670
- [60] Saccà SC, Izzotti A, Rossi P, Traverso C. Glaucomatous outflow pathway and oxidative stress. *Exp Eye Res* 2007; 84(3): 389-99.
<http://dx.doi.org/10.1016/j.exer.2006.10.008> PMID: 17196589
- [61] Saccà SC, Izzotti A. Oxidative stress and glaucoma: Injury in the anterior segment of the eye. *Prog Brain Res* 2008; 173: 385-407.
[http://dx.doi.org/10.1016/S0079-6123\(08\)01127-8](http://dx.doi.org/10.1016/S0079-6123(08)01127-8) PMID: 18929123
- [62] Yokoyama Y, Aizawa N, Chiba N, *et al.* Significant correlations between optic nerve head microcirculation and visual field defects and nerve fiber layer loss in glaucoma patients with myopic glaucomatous disk. *Clin Ophthalmol* 2011; 5: 1721-7.
 PMID: 22205831
- [63] Zhang S, Wu C, Liu L, *et al.* Optical coherence tomography angiography of the peripapillary retina in primary angle-closure glaucoma. *Am J Ophthalmol* 2017; 182: 194-200.
<http://dx.doi.org/10.1016/j.ajo.2017.07.024> PMID: 28797550
- [64] Rao HL, Pradhan ZS, Suh MH, Moghimi S, Mansouri K, Weinreb RN. Optical coherence tomography angiography in glaucoma. *J Glaucoma* 2020; 29(4): 312-21.
<http://dx.doi.org/10.1097/IJG.0000000000001463> PMID: 32053551
- [65] Chetry D, Singh J, Chhetri A, Katiyar VK, Singh DS. Effect of yoga on intra-ocular pressure in patients with glaucoma. *Indian J Ophthalmol* 2023; 71(5): 1757-65.
http://dx.doi.org/10.4103/ijo.IJO_2054_22 PMID: 37203026
- [66] Yadav RK, Sankalp , Dada T, Faiq M. Effect of yoga-based ocular exercises in lowering of intraocular pressure in glaucoma patients: An affirmative proposition. *Int J Yoga* 2018; 11(3): 239-41.
http://dx.doi.org/10.4103/ijoy.IJOY_55_17 PMID: 30233118
- [67] Pandya J, Schardt M, Yu D, Henderer JD. A pilot study on the effect of alternate nostril breathing and foot reflexology on intraocular pressure in ocular hypertension. *J Altern Complement Med* 2019; 25(8): 824-6.
<http://dx.doi.org/10.1089/acm.2019.0033> PMID: 31180234
- [68] Rice R, Allen RC. Yoga in Glaucoma. *Am J Ophthalmol* 1985; 100(5): 738-9.
[http://dx.doi.org/10.1016/0002-9394\(85\)90634-8](http://dx.doi.org/10.1016/0002-9394(85)90634-8) PMID: 4061559
- [69] Udenia H, Mittal S, Agrawal A, Singh A, Singh A, Mittal SK. Yogic pranayama and diaphragmatic breathing: adjunct therapy for

- intraocular pressure in patients with primary open-angle glaucoma: A randomized controlled trial. *J Glaucoma* 2021; 30(2): 115-23.
<http://dx.doi.org/10.1097/IJG.0000000000001697> PMID: 33955942
- [70] Ismail AMA, Abd Elfatah Abo Saif HF, El-Moatasem Mohamed AM. Effect of *Jyoti-Trataka* on intraocular pressure, autonomic control, and blood glucose in diabetic patients with high-tension primary open-angle glaucoma: A randomized-controlled trial. *J Complement Integr Med* 2022; 19(4): 1013-8.
<http://dx.doi.org/10.1515/jcim-2021-0041> PMID: 34303323
- [71] Gagrani M, Faiq MA, Sidhu T, et al. Meditation enhances brain oxygenation, upregulates BDNF and improves quality of life in patients with primary open angle glaucoma: A randomized controlled trial. *Restor Neurol Neurosci* 2018; 36(6): 741-53.
<http://dx.doi.org/10.3233/RNN-180857> PMID: 30400122
- [72] Jasien JV, Jonas JB, de Moraes CG, Ritch R. Intraocular pressure rise in subjects with and without glaucoma during four common yoga positions. *PLoS One* 2015; 10(12): e0144505.
<http://dx.doi.org/10.1371/journal.pone.0144505> PMID: 26698309
- [73] Monteiro de Barros DS, Bazzaz S, Gheith ME, Siam GA, Moster MR. Progressive optic neuropathy in congenital glaucoma associated with the Sirsasana yoga posture. *Ophthalmic Surg Lasers Imaging Retina* 2008; 39(4): 339-40.
<http://dx.doi.org/10.3928/15428877-20080701-03> PMID: 18717444
- [74] Bertschinger DR, Mendrinos E, Dosso A. Yoga can be dangerous glaucomatous visual field defect worsening due to postural yoga. *Br J Ophthalmol* 2007; 91(10): 1413-4.
<http://dx.doi.org/10.1136/bjo.2007.114546> PMID: 17895421
- [75] Zaher O, Kuchtaruk AA, Malvankar-Mehta MS. Effect of various relaxation techniques on the intraocular pressure of patients with glaucoma: Systematic review and meta-analysis. *Can J Ophthalmol* 2023.
PMID: 37364856
- [76] Walker E, Hernandez AV, Kattan MW. Meta-analysis: Its strengths and limitations. *Cleve Clin J Med* 2008; 75(6): 431-9.
<http://dx.doi.org/10.3949/ccjm.75.6.431> PMID: 18595551

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.